

# Spin-1 resonances as a signature of composite Higgs at the LHC

Anna Kamińska

with S.Pokorski, A.Weiler



This research project has been supported by a Marie Curie ITN Fellowship (PITN-GA-2009-237920-UNILHC)

Planck 2013, Bonn, 21.05.13

# QCD inspired

QCD Lagrangian in the limit  $m_u, m_d \rightarrow 0$

$$SU(2)_L \times SU(2)_R \rightarrow SU(2)_V$$

$\sqrt{s} \ll \Lambda_{QCD}$  pions interact weakly  $\rightarrow$  effective description

$$U \rightarrow g_L U g_R^\dagger, \quad U = e^{i\pi\sigma^a/f_\pi}, \quad \mathcal{L}^{(2)} = \frac{f_\pi^2}{4} \text{Tr} \left\{ D^\mu U^\dagger D_\mu U \right\}$$

# QCD inspired

QCD Lagrangian in the limit  $m_u, m_d \rightarrow 0$

$$SU(2)_L \times SU(2)_R \rightarrow SU(2)_V$$

$\sqrt{s} \ll \Lambda_{QCD}$  pions interact weakly  $\rightarrow$  effective description

$$U \rightarrow g_L U g_R^\dagger, \quad U = e^{i\pi\sigma^a/f_\pi}, \quad \mathcal{L}^{(2)} = \frac{f_\pi^2}{4} \text{Tr} \left\{ D^\mu U^\dagger D_\mu U \right\}$$

## SM Higgs sector

$$\mathcal{G} = SU(2)_L \times SU(2)_R \quad \rightarrow \quad \mathcal{H} = SU(2)_C$$

$$\phi = \begin{pmatrix} H^0 & H^+ \\ -H^- & H^0 \end{pmatrix}, \quad \phi \rightarrow g_L \phi g_R^\dagger$$

# QCD inspired

QCD Lagrangian in the limit  $m_U, m_D \rightarrow 0$

$$SU(2)_L \times SU(2)_R \rightarrow SU(2)_V$$

$\sqrt{s} \ll \Lambda_{QCD}$  pions interact weakly  $\rightarrow$  effective description

$$U \rightarrow g_L U g_R^\dagger, \quad U = e^{i\pi\sigma^a/f_\pi}, \quad \mathcal{L}^{(2)} = \frac{f_\pi^2}{4} \text{Tr} \left\{ D^\mu U^\dagger D_\mu U \right\}$$

## SM Higgs sector

$$\mathcal{G} = SU(2)_L \times SU(2)_R \rightarrow \mathcal{H} = SU(2)_C$$

$$\phi = \begin{pmatrix} H^0 & H^+ \\ -H^- & H^0 \end{pmatrix}, \quad \phi \rightarrow g_L \phi g_R^\dagger$$

EW symmetry broken  $\phi = (v + h(x)) e^{i\frac{\pi^a(x)\sigma^a}{v}} = (v + h(x)) U$

$$\text{Tr} \left\{ D_\mu \phi^\dagger D_\mu \phi \right\} = \frac{1}{2} \partial^\mu h \partial_\mu h + \frac{(v + h)^2}{4} \text{Tr} \left\{ D^\mu U^\dagger D_\mu U \right\}$$



# Strong electroweak symmetry breaking

**electroweak symmetry broken by new strong interactions**

composite Higgs - PG boson

## electroweak symmetry broken by new strong interactions

### composite Higgs - PG boson

- $SO(5)/SO(4) \rightarrow 4\pi \rightarrow H$

Minimal Composite Higgs Model  
Agashe, Contino, Pomarol '04

- $SO(6)/SO(5) \rightarrow 5\pi \rightarrow H, a$   
 $SU(4)/Sp(4, C) \rightarrow 5\pi \rightarrow H, s$

Next MCHM  
Gripaios, Pomarol, Riva, Serra '09  
Chacko, Batra '08

- $SO(6)/SO(4) \times SO(2) \rightarrow 8\pi \rightarrow H_1 + H_2$

Minimal Composite Two Higgs Doublets  
Mrazek, Pomarol, Rattazzi, Serra, Wulzer '11

# Signatures of composite Higgs

- modified Higgs couplings

$$\xi = (v/f_\pi)^2$$

- modified Higgs couplings

$$\xi = (v/f_\pi)^2$$

## Minimal Composite Higgs Model

$$SO(5) \rightarrow SO(4) \sim SU(2)_L \times SU(2)_R$$

$$\mathcal{L}^{(2)} = \frac{1}{2} \partial_\mu h \partial^\mu h + \frac{v^2}{4} \left( 1 + 2a \frac{h}{v} + b \frac{h^2}{v^2} + \dots \right) \text{Tr} \left\{ D^\mu U D_\mu U^\dagger \right\}$$

$$a = \sqrt{1 - \xi}, \quad b = 1 - 2\xi.$$



# Signatures of composite Higgs

- modified Higgs couplings

$$\xi = (v/f_\pi)^2$$

## Minimal Composite Higgs Model

$$SO(5) \rightarrow SO(4) \sim SU(2)_L \times SU(2)_R$$

$$\mathcal{L}^{(2)} = \frac{1}{2} \partial_\mu h \partial^\mu h + \frac{v^2}{4} \left( 1 + 2a \frac{h}{v} + b \frac{h^2}{v^2} + \dots \right) \text{Tr} \left\{ D^\mu U D_\mu U^\dagger \right\}$$

$$a = \sqrt{1 - \xi}, \quad b = 1 - 2\xi.$$

→ small values of  $\xi$  preferred,  $\xi \lesssim 0.3$

# Signatures of composite Higgs

indirect (electroweak precision, flavor) and direct effects

- spin-1/2 resonances
- **spin-1 resonances**

Contino, Pappadopulo, Marzocca, Rattazzi

Panico, Wulzer

De Curtis, Redi, Tesi

...

→ analogue of  $\rho$  of QCD

→ KK modes from extra dimension

# Signatures of composite Higgs

indirect (electroweak precision, flavor) and direct effects

- spin-1/2 resonances
- **spin-1 resonances**

Contino, Pappadopulo, Marzocca, Rattazzi  
Panico, Wulzer  
De Curtis, Redi, Tesi

...

→ analogue of  $\rho$  of QCD

→ KK modes from extra dimension

## Goal

- provide a simple, general and self-consistent effective framework to study properties of spin-1 resonances

# Signatures of composite Higgs

indirect (electroweak precision, flavor) and direct effects

- spin-1/2 resonances
- **spin-1 resonances**

Contino, Pappadopulo, Marzocca, Rattazzi  
Panico, Wulzer  
De Curtis, Redi, Tesi  
...

→ analogue of  $\rho$  of QCD

→ KK modes from extra dimension

## Goal

- provide a simple, general and self-consistent effective framework to study properties of spin-1 resonances
- LHC phenomenology

# Signatures of composite Higgs

indirect (electroweak precision, flavor) and direct effects

- spin-1/2 resonances
- **spin-1 resonances**

Contino, Pappadopulo, Marzocca, Rattazzi  
Panico, Wulzer  
De Curtis, Redi, Tesi  
...

→ analogue of  $\rho$  of QCD

→ KK modes from extra dimension

## Goal

- provide a simple, general and self-consistent effective framework to study properties of spin-1 resonances
- LHC phenomenology

## Guideline: QCD

# Effective description of spin-1 resonances

global symmetry breaking  $\mathcal{G} \rightarrow \mathcal{H}$

# Effective description of spin-1 resonances

global symmetry breaking  $\mathcal{G} \rightarrow \mathcal{H}$

## Spin-1 resonances

- in a representation of the unbroken global symmetry of strong dynamics
- 'hidden local symmetry'

global symmetry breaking  $\mathcal{G} \rightarrow \mathcal{H}$

## Spin-1 resonances

- in a representation of the unbroken global symmetry of strong dynamics
  - 'hidden local symmetry'
- modify the symmetry breaking pattern

$$\mathcal{G} \times \mathcal{H}_{local} \rightarrow \mathcal{H}$$

- SM electroweak  $SU(2)_L \times U(1)_Y$  group sits in  $\mathcal{G}$
- gauge bosons of  $\mathcal{H}_{local}$  → 'vector' resonances



# Effective description of spin-1 resonances

global symmetry breaking  $\mathcal{G} \rightarrow \mathcal{H}$

## Spin-1 resonances

- in a representation of the unbroken global symmetry of strong dynamics
  - 'hidden local symmetry'
- modify the symmetry breaking pattern

$$\mathcal{G} \times \mathcal{H}_{local} \rightarrow \mathcal{H}$$

- SM electroweak  $SU(2)_L \times U(1)_Y$  group sits in  $\mathcal{G}$
- gauge bosons of  $\mathcal{H}_{local}$  → 'vector' resonances

$$S \rightarrow g S h^\dagger, \quad g \in \mathcal{G}, \quad h \in \mathcal{H}_{local}, \quad \langle S \rangle = \mathbf{1}.$$

$$\mathcal{L} \ni v_1^2 \text{Tr} \left\{ D_\mu S D^\mu S^\dagger \right\}$$

# Spin-1 resonances

- general features for small  $\xi$ , small  $g/g_\rho$

- eigenstates - mixture of SM and 'hidden gauge' fields

# Spin-1 resonances

- general features for small  $\xi$ , small  $g/g_\rho$

- eigenstates - mixture of SM and 'hidden gauge' fields

at leading order in  $g/g_\rho$

- heavy spin-1 eigenstates  $\leftrightarrow$  'hidden gauge'  $\rho^\mu$  fields
- light eigenstates  $\leftrightarrow$  SM  $A, W, Z$  fields
- mixing  $\sim g, g'/g_\rho \rightarrow$  interactions!


# Spin-1 resonances

- general features for small  $\xi$ , small  $g/g_\rho$

- eigenstates - mixture of SM and 'hidden gauge' fields

at leading order in  $g/g_\rho$

- heavy spin-1 eigenstates  $\leftrightarrow$  'hidden gauge'  $\rho^\mu$  fields
  - light eigenstates  $\leftrightarrow$  SM  $A, W, Z$  fields
  - mixing  $\sim g, g'/g_\rho \rightarrow$  interactions!
- 
- 'pairing up' of  $SU(2)$  subgroups




# Spin-1 resonances

- general features for small  $\xi$ , small  $g/g_\rho$

- eigenstates - mixture of SM and 'hidden gauge' fields

at leading order in  $g/g_\rho$

- heavy spin-1 eigenstates  $\leftrightarrow$  'hidden gauge'  $\rho^\mu$  fields
  - light eigenstates  $\leftrightarrow$  SM  $A, W, Z$  fields
  - mixing  $\sim g, g'/g_\rho \rightarrow$  interactions!
- 
- 'pairing up' of  $SU(2)$  subgroups



- 3 free parameters:  $\xi, g_\rho, g_{\rho\pi\pi}$

$$g_{\rho\pi\pi} \epsilon^{abc} \pi^a \partial_\mu \pi^b \rho_\mu^c - g_\rho \epsilon^{abc} \partial_\mu \rho_\nu^a \rho_\mu^b \rho_\nu^c$$

$$SO(5) \rightarrow SO(4) \sim SU(2)_L \times SU(2)_R$$

- heavy spin-1 eigenstates  $\leftrightarrow \tilde{\rho}_L^\mu, \tilde{\rho}_R^\mu$

# $SO(5) \rightarrow SO(4) \sim SU(2)_L \times SU(2)_R$

- heavy spin-1 eigenstates  $\leftrightarrow \tilde{\rho}_L^\mu, \tilde{\rho}_R^\mu$
- SM gauge fields

$$W_\mu^\pm \approx \tilde{W}_\mu^\pm - \frac{\sqrt{2}}{2} \sqrt{2-\xi} \frac{g}{g_\rho} \tilde{\rho}_{L\mu}^\pm$$

$$Z_\mu \approx \tilde{Z}_\mu - \frac{\sqrt{2-\xi}}{\sqrt{2}} \frac{g^2 - g'^2}{g_\rho \sqrt{g^2 + g'^2}} \tilde{\rho}_{L\mu}^0 - \frac{2\sqrt{2-2\xi}}{(2-\xi)^{3/2}} \frac{g'^2}{g_\rho \sqrt{g^2 + g'^2}} \tilde{\rho}_{R\mu}^0$$

$$A_\mu \approx \tilde{A}_\mu - \sqrt{\frac{2}{2-\xi}} \frac{e}{g_\rho} \tilde{\rho}_{L\mu} + \sqrt{\frac{2-2\xi}{2-\xi}} \frac{e}{g_\rho} \tilde{\rho}_{R\mu}$$

# $SO(5) \rightarrow SO(4) \sim SU(2)_L \times SU(2)_R$

- heavy spin-1 eigenstates  $\leftrightarrow \tilde{\rho}_L^\mu, \tilde{\rho}_R^\mu$
- SM gauge fields

$$W_\mu^\pm \approx \tilde{W}_\mu^\pm - \frac{\sqrt{2}}{2} \sqrt{2-\xi} \frac{g}{g_\rho} \tilde{\rho}_{L\mu}^\pm$$

$$Z_\mu \approx \tilde{Z}_\mu - \frac{\sqrt{2-\xi}}{\sqrt{2}} \frac{g^2 - g'^2}{g_\rho \sqrt{g^2 + g'^2}} \tilde{\rho}_{L\mu}^0 - \frac{2\sqrt{2-2\xi}}{(2-\xi)^{3/2}} \frac{g'^2}{g_\rho \sqrt{g^2 + g'^2}} \tilde{\rho}_{R\mu}^0$$

$$A_\mu \approx \tilde{A}_\mu - \sqrt{\frac{2}{2-\xi}} \frac{e}{g_\rho} \tilde{\rho}_{L\mu} + \sqrt{\frac{2-2\xi}{2-\xi}} \frac{e}{g_\rho} \tilde{\rho}_{R\mu}$$

assumption: couplings of  $\tilde{\rho}$  eigenstates with SM fermions arise only via their admixture in SM  $W_\mu^\pm, Z_\mu$  and  $A_\mu$

- coupling of  $\rho$  to two fermions enhanced for small  $\xi$



# $SO(5) \rightarrow SO(4) \sim SU(2)_L \times SU(2)_R$

- heavy spin-1 eigenstates  $\leftrightarrow \tilde{\rho}_L^\mu, \tilde{\rho}_R^\mu$
- SM gauge fields

$$W_\mu^\pm \approx \tilde{W}_\mu^\pm - \frac{\sqrt{2}}{2} \sqrt{2-\xi} \frac{g}{g_\rho} \tilde{\rho}_{L\mu}^\pm$$

$$Z_\mu \approx \tilde{Z}_\mu - \frac{\sqrt{2-\xi}}{\sqrt{2}} \frac{g^2 - g'^2}{g_\rho \sqrt{g^2 + g'^2}} \tilde{\rho}_{L\mu}^0 - \frac{2\sqrt{2-2\xi}}{(2-\xi)^{3/2}} \frac{g'^2}{g_\rho \sqrt{g^2 + g'^2}} \tilde{\rho}_{R\mu}^0$$

$$A_\mu \approx \tilde{A}_\mu - \sqrt{\frac{2}{2-\xi}} \frac{e}{g_\rho} \tilde{\rho}_{L\mu} + \sqrt{\frac{2-2\xi}{2-\xi}} \frac{e}{g_\rho} \tilde{\rho}_{R\mu}$$

assumption: couplings of  $\tilde{\rho}$  eigenstates with SM fermions arise only via their admixture in SM  $W_\mu^\pm, Z_\mu$  and  $A_\mu$


- coupling of  $\rho$  to two fermions enhanced for small  $\xi$
- coupling of  $\rho$  to two SM gauge bosons suppressed

$$g_{\rho\pi\pi} = \xi \frac{m_\rho^2}{2g_\rho v^2} = \frac{m_\rho^2}{2g_\rho f_\pi^2}$$

# Production and decays

- production dominated by Drell-Yan  $q\bar{q} \rightarrow \rho$


for a specific value of  
 $\xi = 0.2$  and  $g_\rho = 4$



# Production and decays

- production dominated by Drell-Yan  $q\bar{q} \rightarrow \rho$


for a specific value of  $\xi = 0.2$  and  $g_\rho = 4$



- decays mainly to  $hZ$  and  $WW$ , but  $ll$  non-negligible

for a specific value of  $\xi = 0.2$  and  $g_\rho = 4$



$$\Gamma(\rho \rightarrow WW) \sim m_\rho g_\rho^2 \pi$$



# Direct searches

most sensitive: CMS search for dilepton resonances

$$m_\rho^2 \approx \frac{2g_\rho g_{\rho\pi\pi} v^2}{\xi}$$





# Comparison with indirect constraints

most constraining: prediction for  $\hat{S} = \frac{g^2}{16\pi} S$   
assume: saturation of Weinberg sum rules

$\xi = 0.2$

$\xi = 0.07$



# Conclusions

- signatures of composite Higgs - modified Higgs couplings, effects of resonances
- general effective framework for spin-1 resonances → phenomenology
- at small  $\xi$  - the spin-1 resonance coupling to two SM gauge bosons is suppressed, the coupling to two fermions is enhanced
- resonances mainly Drell-Yan produced
- exclusion limits from searches for dilepton resonances, diboson resonances, dijet mass spectra, ...
- the LHC is already probing the parameter space of spin-1 resonances allowed by electroweak precision tests

# Perturbative unitarity constraints

without spin-1 resonances  $\mathcal{M}_{WW \rightarrow WW}^0(s) \sim \frac{1}{16\pi} \frac{\xi s}{v^2} = \frac{1}{16\pi} \frac{s}{f_\pi^2}$


# Perturbative unitarity constraints

without spin-1 resonances  $\mathcal{M}_{WW \rightarrow WW}^0(s) \sim \frac{1}{16\pi} \frac{\xi s}{v^2} = \frac{1}{16\pi} \frac{s}{f_\pi^2}$   
→ perturbative unitarity violation at  $\Lambda \sim 1.3 \text{ TeV}/\sqrt{\xi}$

$$\xi = 0.2$$

add  $\rho_L$  and  $\rho_R$   
resonances,  
inelastic channels  
included

$$m_\rho^2 \approx \frac{2g_\rho g_{\rho\pi\pi} v^2}{\xi}$$






# Effective description of spin-1 resonances

global symmetry breaking  $\mathcal{G} \rightarrow \mathcal{H}$


- 'vector' resonances  $\mathcal{G} \times \mathcal{H}_{local} \rightarrow \mathcal{H}$




# Effective description of spin-1 resonances

global symmetry breaking  $\mathcal{G} \rightarrow \mathcal{H}$

- 'vector' resonances  $\mathcal{G} \times \mathcal{H}_{local} \rightarrow \mathcal{H}$




- 'vector' and 'axial' resonances  $\mathcal{G} \times \mathcal{G}_{local} \rightarrow \mathcal{H}$




# Effective description of spin-1 resonances

global symmetry breaking  $\mathcal{G} \rightarrow \mathcal{H}$

- 'vector' resonances  $\mathcal{G} \times \mathcal{H}_{local} \rightarrow \mathcal{H}$



- 'vector' and 'axial' resonances  $\mathcal{G} \times \mathcal{G}_{local} \rightarrow \mathcal{H}$




- more resonances  $\mathcal{G} \times \mathcal{G}_{local} \times \mathcal{H}_{local} \rightarrow \mathcal{H}$




# Effective description of spin-1 resonances

global symmetry breaking  $\mathcal{G} \rightarrow \mathcal{H}$

- 'vector' resonances  $\mathcal{G} \times \mathcal{H}_{local} \rightarrow \mathcal{H}$



- 'vector' and 'axial' resonances  $\mathcal{G} \times \mathcal{G}_{local} \rightarrow \mathcal{H}$



- more resonances  $\mathcal{G} \times \mathcal{G}_{local} \times \mathcal{H}_{local} \rightarrow \mathcal{H}$



- 'vector' resonances most relevant for phenomenology