

Extreme multiplicity studies in the SVD-2 experiment

E. Kokoulina, A. Kutov, V. Ryadovikov

On behalf of SVD-2 Collaboration (JINR, SINP MSU, JHEP)

Of blessed memory P.F. Ermolov



1932 - 2008












<u>Scientific programme:</u>

multiparticle dynamics study in p+p(A) interactions with the extreme multiplicity (more than mean multiplicity) with 50-70 GeV proton beam on U-70 of IHEP (Protvino).


















SVD-2 setup







Scintillation hodoscope ("camomile") for the registration of rare events with HM:

20 elements ("petals"): triangle h=18, 1.8 mm thick.












SIMULATION:

- ➤ Making of Monte Carlo event generator for modeling of setup elements;
- > Software design of data taking and data processing for HM events;
- ➤ Use of JEANT-3 generator for analysis of program packages of event reconstruction;
- \triangleright the 2006 2008 runs on U-70.










MD of simulated (red) and reconstructed (blue) events for Vertex Detector on X and Y projections








Alignment task: is an essential step in the track reconstruction:



















<u>Sequential histogramming method</u> and also: Kalman filter and others reasonable algorithms for track reconstruction .












MSVD:

- z coordinate of interaction vertex:
- 1) hydrogen target,
- 2) shell and
- 3) hodoscope.









Multiplicity Distributions


(MSVD)










Reconstructed HM - events (examples)












































HM Trigger level determines the bottom boundary of registered multiplicity (n>2, 4, ...)






















DT calibration function



Drift tubes: the efficiency of the registration ~ 99%


Resolution ~200 μm












 $M_{\pi0} = 0.1349 \pm 0.009 \, (GeV)$



Detector of gamma-quantum, 2006 run.







Statistics:

more than $8x10^6$ events and $\sim 6x10^6$ from them with trigger level 6, 7, 8 and 10 were recorded.

We continue data taking and data processing.








GDM:

multiparticle dynamics, predictions of charged and neutral MD, QCD-cascade, hadronization mechanisms.









Base of GDM: We use the unified approach to the multiplicity description at HEP. It is based on essentials of QCD and on the main experimental phenomena in multiparticle dynamics.

We are interested the extreme multiplicity region: $n > \bar{n}(s)$.







GDM for hadron interactions:

- Quarks of initial protons are staying in leading particles. Multiparticle production (MP) is realized by active gluons (behavior of hadronization parameters).
- Recombination mechanism of hadronization.
- Particle production limits.







- The dominance of active gluon plenty in central nuclei collisions leads to the new kind of matter QGP.
- Cluster superposition consisted from one, two or more gluon fission (increasing of MD width).
- ❖ We believe the realization of our program will be successful and unique for MP study.







Search for collective phenomena:


- 1. BEC (condensation).
- 2. Ring events (Cherenkov gluon).
- 3. The excess of soft photon yield.
- 4. Clustering of secondaries.
- 5. Turbulence phenomena.








1) M.I.Gorenstein & V.V.Begun (Phys.Lett.B651:114-118,2007): at HM the experimental manifestation of BEC may be founded by the abrupt and anomalous increase of the scaled variance ω^0 of neutral and charged pion fluctuations in the vicinity of BEC-line.














2) Ring-event searching.







3) Soft Photons:
$$p_t \le 0.1 GeV/c, x \le 0.01$$

 $\sigma(SP)$ are 5-8 times larger than expected ones from QED.

$$\sigma_{\gamma} \approx 4mb$$
, $\sigma_{in} \approx 40mb$, $\sigma_{\gamma} \approx n_{\gamma}(T) \cdot \sigma_{in} \rightarrow n_{\gamma} \approx 0.1$

Assumption: QGS or excited new formed hadrons set in almost equilibrium state during a short period. That is why we use the black body emission spectrum:

$$\frac{dn_{\gamma}}{dv} = \frac{8\pi}{c^3} \frac{v^3}{e^{\frac{hv}{T}} - 1}.$$







Soft photons (p < 50 MeV) excess

The number of SP is estimated by MVB density $\rho(T)=n(T)/V$:

$$n_{\gamma}(T) = 0.244 \cdot V \left(\frac{2\pi kT}{hc}\right)^{3}, T_{r} = 2.725 K(MVB) \rightarrow n_{\gamma}(T_{r})/V = 4.112 \cdot 10^{8} m^{-3}$$

$$T = p \approx p_t \cdot \sqrt{2}$$
 $L^3 \cdot \rho(T) \approx n_{\gamma} \rightarrow L(T)$

- estimation of SP emission region ~ 4-6 fm;
- SP excess is the manifestation of kinematical freeze out.


M.Volkov,E.K., E.Kuraev. Part. & Nucl. Let., 5 (2004)122







4) RUN 2002: the indication to the grouping of secondary in the certain direction (clustering) in pA (Pb). [hep-ph/0612364].















We think on successful accomplishment of our program.

