

# Transverse single-spin asymmetries in $W^\pm$ and $Z^0$ boson production in p+p collisions at $\sqrt{s} = 500$ GeV at RHIC

Monday, 25 August 2014 14:40 (20 minutes)

The Sivers function  $f_{1T}^\perp$  describes the correlation of parton transverse momentum with the transverse spin of the nucleon. There is evidence of a quark Sivers effect in semi-inclusive DIS (SIDIS) measurements.

In SIDIS, the quark Sivers function is associated with a final state effect from the gluon exchange between the struck quark and the target nucleon remnants. On the other hand, for the virtual photon production in the Drell-Yan process, the Sivers asymmetry appears as an initial state interaction effect.

As a consequence, the quark Sivers functions are of opposite sign in SIDIS and in Drell-Yan and this non-universality is a fundamental prediction from the gauge invariance of QCD. The experimental test of this sign change is one of the open questions in hadronic physics, and can provide a direct verification of TMD factorization.

While luminosities required for a precise measurement of asymmetries in Drell-Yan production are challenging,  $W^\pm/Z^0$  production is equally sensitive to the predicted sign change and can be well measured at the STAR experiment. The results can also provide essential input to study the evolution effects of the Sivers function, because of the high  $Q^2$  in  $W^\pm/Z^0$ -production. The  $W$  boson kinematics are fully reconstructed from the decay

lepton and the recoil, by employing a MC-based correction, thus avoiding the dilution that an asymmetry reconstructed from the decay lepton only would suffer. We will present the preliminary results for the measurement of the transverse single spin asymmetry,  $A_N$ , of the  $W^\pm/Z^0$  bosons from the STAR experiment at RHIC using transversely polarized proton-proton collisions at  $\sqrt{s} = 500$  GeV.

RHIC is capable of delivering up to  $\sim 1 \text{ fb}^{-1}$  at  $\sqrt{s} = 500$  GeV during the upcoming 2016 p+p run. This will allow STAR to simultaneously test the Sivers' sign change, investigate the sea-quark Sivers' function and pin down TMD evolution through measurements of Drell-Yan,  $W^\pm/Z^0$  and  $\gamma$  production in p+p collisions.

**Primary author:** Dr FAZIO, Salvatore (Brookhaven National Laboratory)

**Presenter:** Dr FAZIO, Salvatore (Brookhaven National Laboratory)

**Session Classification:** Quarks and gluons in hadrons, the hadron spectrum

**Track Classification:** 2) Quarks and gluons in hadrons, the hadron spectrum