

# Cosmic Axion Spin Precession Experiment (CASPEr)

Surjeet Rajendran,  
Stanford

with

Dmitry Budker,  
Peter Graham,  
Micah Ledbetter,  
Alex Sushkov


**Based On:**

D. Budker et.al.,  
arXiv:1306.6089

P.W. Graham and S.R.,  
PRD 88 (2013) 035023,  
(arXiv:1306.6088)

P.W. Graham and S.R.,  
PRD 84 (2011) 055013  
(arXiv:1101.2691)


# Particle Dark Matter



Non-gravitational interactions

Detect these interactions?

# Dark Matter Candidates



**WIMP**

$M \sim 100 \text{ GeV.}$

Weak interactions.

e.g. Neutralino.

**Ultra-light scalars**

Derivative coupling.

Ultra-high energy  
physics.

e.g. Axions

**Light fermions**

$M \sim \text{keV.}$

High energy physics.

e.g. Gravitino.

(Goodman and Witten, 1985)


Axions and WIMPs are the best motivated cold dark matter candidates

# Axions From High Energy Physics

Easy to generate axions from high energy theories

have a global symmetry broken at a high scale  $f_a$

string theory or extra dimensions naturally  
create axions from non-trivial topology



naturally gives large  $f_a \sim \text{GUT } (10^{16} \text{ GeV})$  or Planck  $(10^{19} \text{ GeV})$  scales

# The QCD Axion


Strong CP problem:

$\mathcal{L} \supset \theta G\tilde{G}$  creates a nucleon EDM  $d \sim 3 \times 10^{-16} \theta \text{ e cm}$

measurements  $\Rightarrow \theta \lesssim 3 \times 10^{-10}$

the axion is a simple solution:

$\mathcal{L} \supset \frac{a}{f_a} G\tilde{G}$  with  $m_a \sim \frac{(200 \text{ MeV})^2}{f_a} \sim \text{MHz} \left( \frac{10^{16} \text{ GeV}}{f_a} \right)$



$$a(t) \sim a_0 \cos(m_a t)$$

cosmic expansion reduces amplitude  $a_0$

this field has momentum = 0  $\Rightarrow$  it is non-relativistic matter


the axion is a good cold dark matter candidate

(J. Preskill et.al. 1982)

# Cosmic Axions

misalignment production:

after inflation axion is a constant field, mass turns on at  $T \sim \Lambda_{\text{QCD}}$  then axion oscillates



$$a(t) \sim a_0 \cos(m_a t)$$

Preskill, Wise & Wilczek, Abbott & Sikivie, Dine & Fischler (1983)

axion easily produces correct abundance  $\rho = \rho_{\text{DM}}$


requires  $\left(\frac{a_i}{f_a}\right) \sqrt{\frac{f_a}{M_{\text{Pl}}}} \sim 10^{-3.5}$  late time entropy production eases this

e.g.  $\frac{f_a}{M_{\text{Pl}}} \sim 10^{-7} \quad \frac{a_i}{f_a} \sim 1 \quad \text{or} \quad \frac{f_a}{M_{\text{Pl}}} \sim 10^{-3} \quad \frac{a_i}{f_a} \sim 10^{-2}$

inflationary cosmology does not prefer flat prior in  $\Theta_i$  over flat in  $f_a$

all  $f_a$  in DM range (all axion masses  $\lesssim \text{meV}$ ) equally reasonable

# Constraints and Searches



# A Different Operator For Axion Detection

So how can we detect high  $f_a$  axions?

Strong CP problem:  $\mathcal{L} \supset \theta G\tilde{G}$  creates a nucleon EDM  $d \sim 3 \times 10^{-16} \theta \text{ e cm}$

the axion:  $\mathcal{L} \supset \frac{a}{f_a} G\tilde{G}$  creates a nucleon EDM  $d \sim 3 \times 10^{-16} \frac{a}{f_a} \text{ e cm}$


$a(t) \sim a_0 \cos(m_a t)$  with  $m_a \sim \frac{(200 \text{ MeV})^2}{f_a} \sim \text{MHz} \left( \frac{10^{16} \text{ GeV}}{f_a} \right)$

axion dark matter  $\rho_{\text{DM}} \sim m_a^2 a^2 \sim (200 \text{ MeV})^4 \left( \frac{a}{f_a} \right)^2 \sim 0.3 \frac{\text{GeV}}{\text{cm}^3}$


so today:  $\left( \frac{a}{f_a} \right) \sim 3 \times 10^{-19}$  independent of  $f_a$

the axion gives all nucleons a rapidly oscillating EDM independent of  $f_a$


# Setup



# Setup



# Outline



1. Signal


2. Parameters

3. Sensitivities

4. Conclusions

# Signal


# Solid State Precision Magnetometry



$$\delta\theta \sim \frac{d_N E}{2\mu_N B - m_a} \sin((2\mu_N B - m_a) t) \sin(2\mu_N B t)$$

Resonant Rabi oscillations

# Solid State Precision Magnetometry



$$\delta B \sim n\mu_N \frac{d_N E}{2\mu_N B - m_a} \sin((2\mu_N B - m_a)t) \sin(2\mu_N Bt)$$

# Rough Estimate

$$\delta B \sim np\mu_N \frac{d_N E}{2\mu_N B - m_a} \sin((2\mu_N B - m_a)t) \sin(2\mu_N Bt)$$

$$n \sim \frac{10^{22}}{\text{cm}^3}$$

$$\mu_N \sim \frac{e}{\text{GeV}}$$

$$d_N \sim 10^{-34} \text{ e-cm}$$

$$p \sim \mathcal{O}(1)$$


$$E_{\text{eff}} \sim 10^6 \frac{\text{V}}{\text{cm}}$$

$$(\mu_N B - m_a)^{-1} \sim (10^{-6} m_a)^{-1} \sim t \sim 1 \text{ s} \left( \frac{f_a}{10^{16} \text{ GeV}} \right)$$

$$\delta B \sim 10^{-2} \text{ fT}$$

# Parameters

# Nuclear Polarization (p)




$$\vec{B}_0 \sim 10 \text{ T}$$

$$\theta_0 \sim 4 \text{ K}$$

$$p \sim \mathcal{O}(1) \text{ for } T_1 \sim 3 \text{ hrs}$$

Optical pumping  
 $p \sim 0.9$  in Liquid Xe, Diamond

# Effective Electric Field




## Schiff's Theorem

$$\text{Equilibrium} \implies \langle \vec{F}_N \rangle = 0$$

Only electrostatic forces on nucleus.

$$\langle \vec{E}_N \rangle = 0$$

# Effective Electric Field



## Schiff Moment


Couple to higher moments of the nucleus.  
(finite size effects)

$$E_{\text{eff}} \sim (10^{-9} Z^3) E_N$$

Use high  $Z$  nucleus (Pb, Hg)

$$E_{\text{eff}} \sim (10^{-3}) E_N$$

# Polar Crystal




## Lead Titanate

Pb displaced from axis of symmetry, large internal field.

$$E_{\text{eff}} \sim 10^6 \frac{\text{V}}{\text{cm}}$$

O. Sushkov et.al.,  
PRA 72, 034501(2005)

# Polar Crystal




$$E_{\text{eff}} \sim 10^6 \frac{\text{V}}{\text{cm}}$$

Field reversal not needed. Ferroelectric not needed.

Larger (~few) fields may exist in other polar crystals.

# Interrogation Time (t)

$$\delta B \sim np\mu_N \frac{d_N E}{2\mu_N B - m_a} \sin((2\mu_N B - m_a)t) \sin(2\mu_N Bt)$$




Temporal coherence of  
the dark matter axion field.

Material limitations.

# Axion Coherence

How large can  $t$  be?




Spatial Homogeneity of the field?

Classical field  $a(x)$  with velocity  $v \sim 10^{-3}$

$$\Rightarrow \frac{\nabla a}{a} \sim m_a v$$

$$t \sim \frac{1}{m_a v^2} = 1 \text{ s} \left( \frac{f_a}{10^{16} \text{ GeV}} \right)$$

# Coherence of Transverse Magnetization



$$B(x_1) \neq B(x_2)$$

Local variations in Larmor frequency leads to dephasing.

$$(T_2)$$

**Spin-Spin Interactions**

$$T_2 \sim 1 \text{ ms}$$

**Dynamic Decoupling**

40 s in Si, 1300 s in Liquid Xe

$$T_2^{\text{eff}} \sim 1 \text{ s}$$

# Recap

$$\delta B \sim np\mu_N \frac{d_N E}{2\mu_N B - m_a} \sin((2\mu_N B - m_a)t) \sin(2\mu_N Bt)$$

$$n \sim \frac{10^{22}}{\text{cm}^3}$$

$$\mu_N \sim \frac{e}{\text{GeV}}$$

$$d_N \sim 10^{-34} \text{ e-cm}$$

$$p \sim \mathcal{O}(1)$$

(e.g. optical pumping)

$$E_{\text{eff}} \sim 10^6 \frac{\text{V}}{\text{cm}}$$

(e.g. polar crystal)

$$(\mu_N B - m_a)^{-1} \sim (10^{-6} m_a)^{-1} \sim t \sim 1 \text{ s} \left( \frac{f_a}{10^{16} \text{ GeV}} \right)$$

(dynamic decoupling and  $m_a > \text{MHz}$ )

$$\delta B \sim 10^{-2} \text{ fT}$$

# Sensitivities

# Noise

1. Magnetization Noise

2. Magnetometer Noise

3. Integration Time

# Magnetization Noise (Spin Projection)



Each spin has random initial transverse projection.

Needs time variation for it to be noise.

$$M_n(\omega) \sim \frac{\mu_N}{r^3} \sqrt{nr^3} \langle S_{\text{rms}}(\omega) \rangle \sim \mu_N \sqrt{\frac{n}{V}} \langle S_{\text{rms}}(\omega) \rangle$$

$$S_{\text{rms}}^2(\omega) \approx \frac{1}{8} \left( \frac{T_2}{1 + T_2^2 (\omega - 2\mu_N B)^2} \right)$$

M. Braun and J. Konig, PRB 75, 085310 (2007)

# Magnetization Noise (Spin Projection)



$$M_n(\omega) \sim \frac{\mu_N}{r^3} \sqrt{nr^3} \langle S_{\text{rms}}(\omega) \rangle \sim \mu_N \sqrt{\frac{n}{V}} \langle S_{\text{rms}}(\omega) \rangle$$


$$S_{\text{rms}}^2(\omega) \approx \frac{1}{8} \left( \frac{T_2}{1 + T_2^2 (\omega - 2\mu_N B)^2} \right)$$

Intuition

$$\omega \gg 2\mu_N B \implies S_{\text{rms}} \sim \sqrt{\frac{1}{T_2 \omega}} \sqrt{\frac{1}{\omega}}$$

$$(\omega - 2\mu_N B) \approx \frac{1}{T_2} \implies S_{\text{rms}} \propto \sqrt{T_2} \text{ (resonantly enhanced)}$$


# Magnetometer Noise (SQUID)



SQUID measures magnetic flux.

More flux with more volume.

# Magnetometer Noise (SQUID)



Typical Parameters

$$\phi_n \sim 10^{-21} \frac{\text{Wb}}{\sqrt{\text{Hz}}} \text{ (at 4 K)}$$

$$L_i \sim 500 \text{ nH} \quad M \sim 10 \text{ nH} \quad r \sim 10 \text{ cm}$$

$$B \sim 0.1 \frac{\text{fT}}{\sqrt{\text{Hz}}}$$

# Integration Time

$$\delta B \sim np\mu_N \frac{d_N E}{2\mu_N B - m_a} \sin((2\mu_N B - m_a)t) \sin(2\mu_N Bt)$$

## Signal to Noise Scaling

$$t \lesssim \min(t_a, T_2) \propto t^{\frac{3}{2}}$$

Signal builds linearly. Noise integrates down.

$$T_2 \lesssim t \lesssim t_a \propto \sqrt{t}$$

Signal limited by  $T_2$ . Noise integrates down.

$$t \gtrsim t_a \propto t^{\frac{1}{4}}$$

Signal looks like excess noise.

$$\sqrt{\rho_n^2 + \rho_a^2} \cong \rho_n + \frac{1}{2} \frac{\rho_a^2}{\rho_n} \sim \frac{\rho_n}{\sqrt{t}} + \frac{\rho_a^2}{2\rho_n}$$

# Integration Time

$$\delta B \sim np\mu_N \frac{d_N E}{2\mu_N B - m_a} \sin((2\mu_N B - m_a)t) \sin(2\mu_N Bt)$$

## Signal to Noise Scaling

$$t \lesssim \min(t_a, T_2) \propto t^{\frac{3}{2}}$$

Signal builds linearly. Noise integrates down.

$$T_2 \lesssim t \lesssim t_a \propto \sqrt{t}$$


Signal limited by  $T_2$ . Noise integrates down.

$$t \gtrsim t_a \propto t^{\frac{1}{4}}$$

Signal looks like excess noise.

Limited by time needed to scan over the full band.

# Projected Sensitivity in Lead Titanate




$$\delta B = 0.1 \frac{\text{fT}}{\sqrt{\text{Hz}}}, n = \frac{10^{22}}{\text{cm}^3}, V = 1000 \text{ cm}^3, T_2 = 1 \text{ s}$$

# Another Operator


$$\mathcal{L} \supset \frac{\partial_\mu a}{f_a} \bar{\psi} \gamma^\mu \gamma_5 \psi$$

$$H \supset \frac{a}{f_a} m_a \vec{v} \cdot \vec{S}$$



Can use Xe, He...

# Projected Sensitivity



$$\delta B = 0.1 \frac{\text{fT}}{\sqrt{\text{Hz}}}, \quad n = \frac{10^{22}}{\text{cm}^3}, \quad V = 1000 \text{ cm}^3, \quad T_2 = 1 \text{ s}$$

Optimal Polar Crystal



Longer  $T_2$



Future



Volume



Magnetometry  
(e.g. atomic magnetometers)

# Conclusions


# Morals

1. Model independent, non-derivative coupling. Phase measurement.  
Moderate scaling with  $f_a$ .
2. Signal  $\propto \sqrt{\rho}$ , can search for small component of dark matter.
3. A/C signal. Resonant boost. Noise amelioration.
4. Signal verification using dependence on electric field and spatial coherence of the axion. Help reject technical noise.
5. Scan over wide bandwidth by changing magnetic field.
6. Scalable. Complements solid state EDM efforts.

# WIMP Detection


Goodman and Witten, 1985 :  $\sigma_{\chi N} \sim 10^{-38} \text{ cm}^2$

Today




Axions deserve similar effort.

# Solid State Axion Searches



# Backup

# Nuclear Polarization (p)




$$\vec{B}_0 \sim 10 \text{ T}$$

$$\theta_0 \sim 4 \text{ K}$$

$$p \sim 10^{-3} \text{ in } T_1 \sim 3 \text{ hrs}$$

# Optical Pumping



$$\vec{B}_0 \sim 10 \text{ T}$$

$$\theta_0 \sim 4 \text{ K}$$

Polarize impurity.


Transfer to nuclei through optical interactions.

# Dynamic Decoupling

$$H(\vec{S}_i) = \alpha_0 + \alpha_1 t + \alpha_2 t^2 + \dots$$

Refocus spins within  $T_2$  through EM pulse sequences.

## Hahn Echo




$\pi$  rotation eliminates unknown but constant gradient.

# Axion-like Particles (ALPs)

Broken global symmetry couples to Standard model through derivative interactions of the Goldstone boson.

Interactions:  $\frac{\partial_\mu a}{f_a} \bar{\Psi} \gamma^\mu \gamma^5 \Psi, \frac{a}{f_a} F^{\mu\nu} \tilde{F}_{\mu\nu}$

Mass:  $m_a$



$$a(t) \sim a_0 \cos(m_a t)$$

cosmic expansion reduces amplitude  $a_0$


this field has momentum = 0  $\Rightarrow$  it is non-relativistic matter

Good cold dark matter candidate

# Axion-like Particles

$$\mathcal{L} \supset \frac{\partial_\mu a}{f_a} \bar{N} \gamma^\mu \gamma^5 N \implies \frac{\langle a \rangle m_a}{f_a} \vec{v} \cdot \vec{S}_N$$

Spin precession perpendicular to galactic dark matter wind.



Electric field/Schiff moment unimportant. Can use low Z.

# Dynamic Decoupling

$$H(\vec{S}_i) = \alpha_0 + \alpha_1 t + \alpha_2 t^2 + \dots$$

Refocus spins within  $T_2$  through EM pulse sequences.

In principle,  $T_2^{\text{eff}} \sim T_1 \sim \text{hr}$

## Demonstrated

(1) 40 s in  $^{29}\text{Si}$  (Y. Dong et.al., PRL 100, 247601 (2008))

(2) 1300 s in Xe (M. Ledbetter and M. Romalis)

$$T_2^{\text{eff}} \sim 1 \text{ s}$$

# Magnetization Noise (Spin Projection)



$$M_n(\omega) \sim \frac{\mu_N}{r^3} \sqrt{nr^3} \langle S_{\text{rms}}(\omega) \rangle \sim \mu_N \sqrt{\frac{n}{V}} \langle S_{\text{rms}}(\omega) \rangle$$

$$\frac{dS}{dt} = -\frac{S}{T_2} + 2\mu_N B \times S$$

$$S_{\text{rms}}^2(\omega) \approx \frac{1}{8} \left( \frac{T_2}{1 + T_2^2 (\omega - 2\mu_N B)^2} \right)$$

M. Braun and J. Konig, PRB 75, 085310 (2007)