

Dark radiation in the non-sequestered Large Volume scenario

in collab. with **P. Mangat, F. Rompineve and L. Witkowski**
(Heidelberg)

Outline

- Dark radiation – observational hints/bounds
- Dark radiation in the sequestered LVS
- ...and in the non-sequestered LVS
- ...and in the LVS with flavor-branes
- Summary/Conclusions

Introduction

- conventional variable: N_{eff}

(effective number of neutrino species; $N_{\text{eff}}^{\text{SM}} = 3.046$)

- Planck + WMAP + highL + BAO:

$$N_{\text{eff}} = 3.3 \pm 0.5 \quad (95\% \text{ CL})$$

- Including also H_0 :

$$N_{\text{eff}} = 3.5 \pm 0.5 \quad (95\% \text{ CL})$$

- \Rightarrow mild preference for $\Delta N_{\text{eff}} \neq 0$;

Here: View this as a bound on dark radiation

- Crucial: Significant improvement expected in the future;
Potential to exclude models with $\Delta N_{\text{eff}} \neq 0$

Introduction - continued

- Conventional picture of cosmological evolution with some **extra light d.o.f. (DR)** :


Inflaton \longrightarrow (Modulus Φ) \longrightarrow SM + DR

$$\Delta N_{eff} = \frac{43}{7} \left(\frac{10.75}{g_*(T_d)} \right)^{1/3} \frac{\rho_{DR}}{\rho_{SM}} \Big|_{T_d}$$

- Here T_d is the decay temperature of Φ and

$$\frac{\rho_{DR}}{\rho_{SM}} \Big|_{T_d} = \frac{\Gamma_{\Phi \rightarrow DR}}{\Gamma_{\Phi \rightarrow SM}}$$

Dark radiation in the Large Volume scenario



- Notation: $T_b = \tau_b + ia_b$; $T_s = \tau_s + ia_s$

$$K = -2 \ln \mathcal{V} = -2 \ln \left((T_b + \bar{T}_b)^{3/2} - (T_s + \bar{T}_s)^{3/2} \right)$$

- Crucial point:
 α' -corrections + non-pert. effects lead to stabilization at exponentially large volume


$$\tau_b \sim \exp(\tau_s) \sim \exp(-\chi)$$

- Classical shift symmetry $a_b \rightarrow a_b + \text{const.}$ is only broken non-perturbatively; $m_a = 0$ for all practical purposes.

Dark radiation in the sequestered Large Volume scenario

Cicoli, Conlon, Quevedo '12

Higaki, Nakayama, Takahashi '12... '13



- SM on fract. D3s at singularity of type-IIB CY-orientifold
- gauge-kinetic function $f = f(S)$
- sequestered Kähler potential:

$$K = -3 \ln \left(T_b + \overline{T}_b - \left[Q^i \overline{Q}^i + H_u \overline{H}_u + \{z H_u H_d + \text{h.c.}\} + \dots \right] \right)$$

see e.g. Blumenhagen, Conlon, Krippendorf, Moster, Quevedo, '09

- A straightforward analysis gives:

$$\Gamma_{\Phi \rightarrow a_b a_b} = \frac{1}{48\pi} \frac{m_\Phi^3}{M_P^2}$$

$$\Gamma_{\Phi \rightarrow H_u H_d} = \frac{2z^2}{48\pi} \frac{m_\Phi^3}{M_P^2}$$

- Conclusion: Need either $z > 2$ or $n_H > 4$.

(Here n_H counts pairs of Higgs doublets and one assumes the bound $N_{eff} < 4$.)

- Comment: Shift symmetry singles out $z = 1$,


$$K_H \sim |H_u + \overline{H}_d|^2.$$

Brignole et al. '95; Choi et al. '03; Bruemmer et al. '08-10;
Ben-Dayan, Einhorn '10; AH, Knochel, Weigand '11

(It is unclear how to realize $z \gg 1$ at a fundamental level.
Note that the Kähler metric becomes singular in this limit.)

Dark radiation in the non-sequestered Large Volume scenario

- The non-sequestered case has been discussed before (in a racetrack-LVS hybrid, with SM on non-pert. stablized cycles)
Higaki, Kamada, Takahashi '12
- It is claimed that axions are not an issue **at all**, but stringy realizability of this specific setting is unclear
- We focus on the (in our opinion more standard)
D-term stabilization of 4-cycle ratios
- We assume that τ_{SM}/τ_b is stabilized by $V_D = 0$.



- Due to SUSY, we then have $T_{SM} = \alpha T_b$, with $\alpha \ll 1$ to be realized by the tuning of gauge fluxes.
- Now $m_{soft} \sim 1/\mathcal{V}$, while $m_{T_b} \sim 1/\mathcal{V}^{3/2}$.

(Thus, low-scale SUSY is difficult to realize cosmologically. But this may actually be OK nowadays...)

- The gauge kinetic function reads

$$f = T_{SM} + hS = \alpha T_b + hS$$

- Again, a straightforward analysis gives:

$$\Gamma_{\Phi \rightarrow a_b a_b} = \frac{1}{48\pi} \frac{m_\Phi^3}{M_P^2}$$

$$\Gamma_{\Phi \rightarrow hh} = \frac{z^2 \sin^2(2\beta)}{192\pi} \frac{m_\Phi^3}{M_P^2}$$

$$\Gamma_{\Phi \rightarrow AA} = \frac{N_g \gamma^2}{96\pi} \frac{m_\Phi^3}{M_P^2}$$

where

$$\gamma = \frac{\tau_{SM}}{\tau_{SM} + h \operatorname{Re} S}$$

- The branching ratio to axions is

$$B_a = \frac{\Gamma_{aa}}{\Gamma_{aa} + \Gamma_{hh} + \Gamma_{AA}} = \frac{1}{1 + \frac{\sin^2(2\beta)}{4} z^2 + \frac{N_g}{2} \gamma^2}$$

- This gives

$$\Delta N_{eff} = \frac{43}{7} \left(\frac{10.75}{g_*(T_d)} \right)^{1/3} \frac{B_a}{1 - B_a}$$


- Thus, assuming $\tan \beta = 1$ and $z \lesssim 1$
and taking $N_g = 12$ (SM),
our only option for lowering B_a is to increase γ .

ΔN_{eff}

in

non-sequestered


LVS



Non-sequestered LVS, stabilization by loop corrections

- Known possibility: fibre inflation

Cicoli, Burgess, Quevedo '08



$$\mathcal{V} = \sqrt{\tau_1 \tau_2} - \tau_s^{3/2}$$

- Here $\mathcal{V}_{K3} = \tau_1$; $\mathcal{V}_{Base} = \frac{\tau_2}{\sqrt{\tau_1}}$
- Loops and standard LVS naturally stabilize $\tau_2 \gg \tau_1 \gg \tau_s$.

- Here, the overall volume \mathcal{V} is **not** the lightest modulus
- This role is taken over by the ratio τ_2/τ_1
- Advantage: $\tau_{SM} \sim \tau_1$ is naturally much smaller than the typical volume size.
- Now we have two axions (from T_1 and T_2),

$$B_a = \frac{1}{1 + \frac{1}{5}z^2 + \frac{24}{5}\gamma^2}$$

(for $\tan \beta = 1$ and $N_g = 12$)

- Numerical results are similar to the ‘ D -term case’ above

Fundamental problem:

- In both case, unavoidably $\mathcal{L} \supset T_{light} W_{\alpha}^{SM} W^{SM,\alpha} \Big|_{\theta^2}$
- Our light axion is **also** the QCD axion,
with f_a typically too large
- Way out: Increase \mathcal{V}
(But this lowers $T_{reh.}$ and makes baryogenesis difficult)
- Way out: Accept fine-tuning $a_{initial} \ll a_{typical}$
(This can be justified e.g. if ρ_{DM} is anthropically bounded)


see e.g. Hertzberg, Tegmark, Wilczek '08; Freivogel '08

- Way out: Add a field-theoretic (open-string) QCD axion,
with a decay constant which is set by some
field-theory VEV (\ll string scale)

Yet another possibility:


“sequestered” (or “de-sequestered”) LVS with **flavor branes**

...appearing already in Aldazabal, Ibanez, Quevedo, Uranga '00



- The SM is again at a singularity, but an extra weakly coupled gauge theory lives on a stack of **flavor branes**.
- This gauge theory must be spontaneously broken (Z' bounds apply)
- Cosmology: $\Phi \rightarrow DR + A'_\mu$; Subsequently $A'_\mu \rightarrow SM$
- Second decay is fast; The analysis is (essentially) as before

ΔN_{eff}
in LVS
with
flavor-branes



Conclusions / Summary

- Interpreting present 'dark radiation data' as bounds, the sequestered LVS may already be in trouble

(Although this depends on $T_{reh.}$)
- The 'non-sequestered' or 'de-sequestered' (through flavor branes) LVS provide a natural ways out
- Nevertheless, discovery of dark radiation is expected in the foreseeable future
- Otherwise, there is the potential of ruling out large parts of the LVS parameter space altogether

(Unless one is prepared to accept an anthropically unmotivated tuning)