

Tagged spectator DIS off a polarized spin-1 target

Wim Cosyn


Ghent University, Belgium

DIS 2016
DESY Hamburg

in collaboration with
Ch. Weiss (JLab) & M. Sargsian (FIU)



Tagged spectator DIS process with deuteron



- DIS off a nuclear target with a slow (relative to nucleus c.m.) nucleon detected in the final state
- Control nuclear configuration
- Advantages for the deuteron
 - ▶ simple NN system, non-nucleonic ($\Delta\Delta$) dof suppressed
 - ▶ active nucleon identified
 - ▶ recoil momentum selects nuclear configuration (medium modifications)
 - ▶ limited possibilities for nuclear FSI, calculable
- Wealth of possibilities to study (nuclear) QCD dynamics
- Will be possible in a wide kinematic range @ EIC (**polarized** for JLEIC)
 - ▶ dedicated talk Wed. in WG7

What is needed?

- General expression of SIDIS for a polarized spin 1 target
 - ▶ Tagged spectator DIS is SIDIS in the target fragmentation region

$$\vec{e} + \vec{T} \rightarrow e' + X + h$$

- Dynamical model to express structure functions of the reaction
 - ▶ First step: impulse approximation (IA) model
- Light-front structure of the deuteron
 - ▶ Natural for high-energy reactions as **off-shellness of nucleons** in LF quantization remains **finite**

Polarized spin 1 particle

- Spin state described by a 3*3 density matrix in a basis of spin 1 states polarized along the collinear virtual photon-target axis

$$W_D^{\mu\nu} = \text{Tr}[\rho_{\lambda\lambda'} W^{\mu\nu}(\lambda'\lambda)]$$

- Characterized by **3 vector** and **5 tensor** parameters

$$\mathbf{S}^\mu = \langle \hat{W}^\mu \rangle, \quad \mathbf{T}^{\mu\nu} = \frac{1}{2} \sqrt{\frac{2}{3}} \langle \hat{W}^\mu \hat{W}^\nu + \hat{W}^\nu \hat{W}^\mu + \frac{4}{3} \left(g^{\mu\nu} - \frac{\hat{P}^\mu \hat{P}^\nu}{M^2} \right) \rangle$$

- Split in longitudinal and orthogonal components

$$\rho_{\lambda\lambda'} = \frac{1}{3} \begin{bmatrix} 1 - \frac{3}{2} S_L + \sqrt{\frac{3}{2}} T_{LL} & \frac{3}{2\sqrt{2}} S_\perp e^{+i(\phi_h - \phi_S)} & \sqrt{\frac{3}{2}} T_{\perp\perp} e^{+i(2\phi_h - 2\phi_T)} \\ & + \sqrt{3} T_{L\perp} e^{+i(\phi_h - \phi_T)} & \\ \frac{3}{2\sqrt{2}} S_\perp e^{-i(\phi_h - \phi_S)} & 1 - \sqrt{6} T_{LL} & \frac{3}{2\sqrt{2}} S_\perp e^{i(\phi_h - \phi_S)} \\ + \sqrt{3} T_{L\perp} e^{-i(\phi_h - \phi_T)} & & - \sqrt{3} T_{L\perp} e^{i(\phi_h - \phi_T)} \\ \sqrt{\frac{3}{2}} T_{\perp\perp} e^{-i(2\phi_h - 2\phi_T)} & - \frac{3}{2\sqrt{2}} S_\perp e^{-i(\phi_h - \phi_S)} & 1 + \frac{3}{2} S_L + \sqrt{\frac{3}{2}} T_{LL} \\ & - \sqrt{3} T_{L\perp} e^{-i(\phi_h - \phi_T)} & \end{bmatrix}$$

Spin 1 SIDIS: General structure of cross section

- To obtain structure functions, enumerate all possible tensor structures that obey hermiticity and QED Ward identity
- Cross section has 41 structure functions,

$$\frac{d\sigma}{dx dQ^2 d\phi_{l'}} = \frac{y^2 \alpha^2}{Q^4(1-\epsilon)} (F_U + F_S + F_T) d\Gamma_{P_h},$$

- ▶ U + S part identical to spin 1/2 case [Bacchetta et al. JHEP ('07)]

$$F_U = F_{UU,T} + \epsilon F_{UU,L} + \sqrt{2\epsilon(1+\epsilon)} \cos \phi_h F_{UU}^{\cos \phi_h} + \epsilon \cos 2\phi_h F_{UU}^{\cos 2\phi_h} + h \sqrt{2\epsilon(1-\epsilon)} \sin \phi_h F_{LU}^{\sin \phi_h}$$

$$\begin{aligned} F_S = & \textcolor{red}{S_L} \left[\sqrt{2\epsilon(1+\epsilon)} \sin \phi_h F_{US_L}^{\sin \phi_h} + \epsilon \sin 2\phi_h F_{US_L}^{\sin 2\phi_h} \right] \\ & + \textcolor{red}{S_L} \textcolor{brown}{h} \left[\sqrt{1-\epsilon^2} F_{LS_L} + \sqrt{2\epsilon(1-\epsilon)} \cos \phi_h F_{LS_L}^{\cos \phi_h} \right] \\ & + \textcolor{red}{S_L} \left[\sin(\phi_h - \phi_S) \left(F_{US_{T,T}}^{\sin(\phi_h - \phi_S)} + \epsilon F_{US_{T,L}}^{\sin(\phi_h - \phi_S)} \right) + \epsilon \sin(\phi_h + \phi_S) F_{US_T}^{\sin(\phi_h + \phi_S)} \right. \\ & \left. + \epsilon \sin(3\phi_h - \phi_S) F_{US_T}^{\sin(3\phi_h - \phi_S)} + \sqrt{2\epsilon(1+\epsilon)} \left(\sin \phi_S F_{US_T}^{\sin \phi_S} + \sin(2\phi_h - \phi_S) F_{US_T}^{\sin(2\phi_h - \phi_S)} \right) \right] \\ & + \textcolor{red}{S_L} \textcolor{brown}{h} \left[\sqrt{1-\epsilon^2} \cos(\phi_h - \phi_S) F_{LS_T}^{\cos(\phi_h - \phi_S)} + \right. \\ & \left. \sqrt{2\epsilon(1-\epsilon)} \left(\cos \phi_S F_{LS_T}^{\cos \phi_S} + \cos(2\phi_h - \phi_S) F_{LS_T}^{\cos(2\phi_h - \phi_S)} \right) \right], \end{aligned}$$

Spin 1 SIDIS: General structure of cross section


- To obtain structure functions, enumerate all possible tensor structures that obey hermiticity and QED Ward identity
- Cross section has 41 structure functions,

$$\frac{d\sigma}{dx dQ^2 d\phi_{l'}} = \frac{y^2 \alpha^2}{Q^4(1-\epsilon)} (F_U + F_S + F_T) d\Gamma_{P_h},$$

- ▶ 23 SF unique to the spin 1 case (tensor pol.), 4 survive in inclusive (b_{1-4})

$$\begin{aligned} F_T = & T_{LL} \left[F_{UT_{LL},T} + \epsilon F_{UT_{LL},L} + \sqrt{2\epsilon(1+\epsilon)} \cos \phi_h F_{UT_{LL}}^{\cos \phi_h} + \epsilon \cos 2\phi_h F_{UT_{LL}}^{\cos 2\phi_h} \right] \\ & + T_{LL} h \sqrt{2\epsilon(1-\epsilon)} \sin \phi_h F_{LT_{LL}}^{\sin \phi_h} \\ & + T_{L\perp} [\dots] + T_{L\perp} h [\dots] \\ & + T_{\perp\perp} \left[\cos(2\phi_h - 2\phi_{T_\perp}) \left(F_{UT_{TT},T}^{\cos(2\phi_h - 2\phi_{T_\perp})} + \epsilon F_{UT_{TT},L}^{\cos(2\phi_h - 2\phi_{T_\perp})} \right) \right. \\ & \left. + \epsilon \cos 2\phi_{T_\perp} F_{UT_{TT}}^{\cos 2\phi_{T_\perp}} + \epsilon \cos(4\phi_h - 2\phi_{T_\perp}) F_{UT_{TT}}^{\cos(4\phi_h - 2\phi_{T_\perp})} \right. \\ & \left. + \sqrt{2\epsilon(1+\epsilon)} \left(\cos(\phi_h - 2\phi_{T_\perp}) F_{UT_{TT}}^{\cos(\phi_h - 2\phi_{T_\perp})} + \cos(3\phi_h - 2\phi_{T_\perp}) F_{UT_{TT}}^{\cos(3\phi_h - 2\phi_{T_\perp})} \right) \right] \\ & + T_{\perp\perp} h [\dots] \end{aligned}$$

Tagged DIS with deuteron: model for the IA




- Hadronic tensor can be written as a product of nucleon hadronic tensor with deuteron light-front densities

$$W_D^{\mu\nu}(\lambda', \lambda) = 4(2\pi)^3 \frac{\alpha_R}{2 - \alpha_R} \sum_{i=U,z,x,y} W_{N,i}^{\mu\nu} \rho_D^i(\lambda', \lambda),$$

- Nucleon hadr tensor has standard (un)polarized contributions
 - Effective Bjorken \tilde{x} depends on recoil momentum ($\alpha_R, \mathbf{p}_{R\perp}$)
- $W_{N,U}^{\mu\nu} = -F_{1N}(g^{\mu\nu} + e_q^\mu e_q^\nu) + F_{2N} \frac{L_n^\mu L_n^\nu}{(p_n q)}$ $W_{N,i}^{\mu\nu} = -i \epsilon^{\mu\nu\rho\sigma} \frac{m_N q_\rho}{(p_i q)} \left[s_{i,\sigma} (\mathbf{g}_{1N} + \mathbf{g}_{2N}) - \frac{(qs_i)}{(p_n q)} p_{n,\sigma} \mathbf{g}_{2N} \right]$
- $\rho_D^U(\lambda', \lambda)$ related distribution of **unpolarized** nucleons in the deuteron
- $\rho_D^z(\lambda', \lambda)$ to **longitudinally** pol. nucleon distribution (deut. "helicity")
- $\rho_D^{x,y}(\lambda', \lambda)$ to **transversally** pol. nucleon distr. (deut. "transversity")

Tagged DIS with deuteron: model for the IA



- Hadronic tensor can be written as a product of nucleon hadronic tensor with deuteron light-front densities


$$W_D^{\mu\nu}(\lambda', \lambda) = 4(2\pi)^3 \frac{\alpha_R}{2 - \alpha_R} \sum_{i=U,z,x,y} W_{N,i}^{\mu\nu} \rho_D^i(\lambda', \lambda),$$

Allows us to write all of the 41 conditional structure functions as a product of a factor dep. on nucleon SF and a D spectral function dep. on the polarization state.
One example:

$$F_{UU}^{\cos 2\phi_h} = \frac{|\mathbf{p}_{R\perp}|}{(p_n q)} F_{2N}(x, \alpha_R, \mathbf{p}_{R\perp}) \times [U(k)^2 + W(k)^2] \frac{(2\pi)^3 E_k}{\pi(2 - \alpha_R)^2}.$$

- In the IA the following structure functions are **zero**
 - lepton polarized single-spin asymmetry [$F_{LU}^{\sin \phi_h}$]
 - target vector polarized single-spin asymmetry [8 SFs]
 - target tensor polarized double-spin asymmetry [7 SFs]

Deuteron light-front wave function




- Up to momenta of a few 100 MeV dominated by NN component
- Can be evaluated in LFQM [Coester,Keister,Polyzou et al.] or covariant Feynman diagrammatic way [Frankfurt,Sargsian,Strikman]
- One obtains a Schrödinger (non-rel) like eq. for the wave function components
- Light-front WF obeys baryon and momentum sum rule

$$\Psi_{\lambda}^D(\mathbf{k}_f, \lambda_1, \lambda_2) = \sqrt{E_{kf}} \sum_{\lambda'_1 \lambda'_2} \mathcal{D}_{\lambda'_1 \lambda'_2}^{\frac{1}{2}} [\mathcal{R}_{fc}(k_{1_f}^{\mu} / m_N)] \mathcal{D}_{\lambda'_2 \lambda'_2}^{\frac{1}{2}} [\mathcal{R}_{fc}(k_{2_f}^{\mu} / m_N)] \Phi_{\lambda}^D(\mathbf{k}_f, \lambda'_1, \lambda'_2)$$


- Differences with non-rel wave function:
 - ▶ appearance of the **Melosh rotations** to account for light-front quantized nucleon states
 - ▶ 3-momentum \mathbf{k}_f is the relative momentum of the nucleons in the light-front boosted rest-frame of the free 2-nucleon state (so not a "true" kinematical variable)

Pole extrapolation for on-shell nucleon structure




- Allows to extract free neutron structure
 - ▶ Recoil momentum p_R controls off-shellness of neutron $t - m_N^2$
 - ▶ Free neutron at pole $t - m_N^2 \rightarrow 0$: “on-shell extrapolation”
 - ▶ Small deuteron binding energy results in small extrapolation length
 - ▶ Eliminates nuclear binding and FSI effects [Sargsian, Strikman PLB '05]
- D-wave suppressed at on-shell point \rightarrow neutron $\sim 100\%$ polarized
- Precise measurements of neutron structure at an EIC

Unpolarized structure function



- Extrapolation for $(m_N^2 - t) \rightarrow 0$ corresponds to on-shell neutron $F_{2N}(x, Q^2)$
- Clear effect of deuteron D-wave, largest in the region dominated by the tensor part of the NN -interaction
- D-wave drops out at the on-shell point

Polarized structure function



- Spin asymmetry $A_{\parallel} = \frac{\sigma(++) - \sigma(-+)}{\sigma(++) + \sigma(-+)} = \frac{F_{LS_L}}{F_T + \epsilon F_L} \propto \frac{g_{1n}}{F_{1n}}$
- Again clear contribution from D-wave at finite recoil momenta
- Relativistic nuclear effects through Melosh rotations, grow with recoil momenta
- Both effects drop out near the on-shell extrapolation point

Outlook

- Study of tensor polarized observables
- Final-state interactions, shadowing corrections
- Medium modifications at higher nucleon momenta
- More exclusive final-states (SIDIS in neutron current fragmentation region)

Conclusion

- General form of SIDIS with a spin 1 target, 23 tensor polarized structure functions unique to spin 1
- Results for the impulse approximation using deuteron light-front structure
- Important contributions from deuteron D -wave, Melosh rotations at larger spectator momenta. Become small if one does pole extrapolation of observables.
- Range of applications at an EIC (with polarized deuterons)