

Feynman integrals, L-series and Kloosterman moments

David Broadhurst, Open University, UK
DESY-Zeuthen, 23 October 2017

I consider Feynman integrals from equal-mass **sunrise** diagrams in two dimensions. At 2 and 3 loops, **elliptic polylogarithms** are obtained from **eta quotients** for **modular forms**. New modular forms appear, on-shell, at 4 and 6 loops. **Kloosterman** moments define **L-series** for all loops, with **rational** relations found up to 22 loops. For all loops, **quadratic** relations between integrals are encoded by **Betti** and **de Rham** matrices.

1. **Honeycomb** walks at **2** loops with **Gauss**
2. **Diamond** walks at **3** loops with **Bessel**
3. **Modular forms** to **6** loops and a **challenge** for RISC
4. **L-series** up to **22** loops
5. **Betti** and **de Rham** matrices for **all** loops

Joint work with David P. Roberts, University of Minnesota Morris, USA

1 Honeycomb walks at 2 loops with Gauss

Let $W_3(2k)$ be the number of returning walks of length $2k$ on a **honeycomb**. In 1960, **Cyril Domb** (1920–2012) showed that

$$G_3(y) = \sum_{k=0}^{\infty} W_3(2k)y^{2k} = 1 + 3y^2 + 15y^4 + 93y^6 + 639y^8 + 4653y^{10} + O(y^{12})$$

is the reciprocal of an **arithmetic-geometric mean** (AGM).

For positive real (a_0, b_0) , Gauss evaluated the **elliptic** integral

$$\frac{2}{\pi} \int_0^{\pi/2} \frac{d\theta}{\sqrt{(a_0 \sin \theta)^2 + (b_0 \cos \theta)^2}} = \frac{1}{\text{agm}(a_0, b_0)}$$

by the rapidly converging process of his AGM:

$$a_{n+1} = \frac{a_n + b_n}{2}, \quad b_{n+1} = \sqrt{a_n b_n}, \quad \text{agm}(a_0, b_0) \equiv a_{\infty} = b_{\infty}.$$

The honeycomb problem is solved by

$$G_3(y) = \frac{1}{\text{agm}(\sqrt{(1-y)^3(1+3y)}, \sqrt{(1+y)^3(1-3y)})}.$$

1.1 Two-loop sunrise diagram with Bessel and Gauss

The two-loop massive **sunrise** diagram in two spacetime dimensions, with external energy w and internal masses (a, b, c) , gives the **Bessel** moment

$$\begin{aligned} I(w, a, b, c) &= 4 \int_0^\infty I_0(wt) K_0(at) K_0(bt) K_0(ct) t dt \\ &= \int_0^\infty \int_0^\infty \frac{dx dy}{P(x, y, 1)} \end{aligned}$$

with $P(x, y, z) = (a^2x + b^2y + c^2z)(xy + yz + zx) - w^2xyz$ obtained from Schwinger parameters. Bailey, Borwein, Broadhurst and Glasser obtained

$$I(w, a, b, c) = 8\pi \int_{a+b+c}^\infty \frac{A(v) v dv}{v^2 - w^2},$$

$$A(w) = 1/\text{agm} \left(\sqrt{F(w)}, \sqrt{16abcw} \right),$$

$$F(w) \equiv (w + a + b + c)(w + a - b - c)(w - a + b - c)(w - a - b + c).$$

Since $F(w) - 16abcw = F(-w)$, the complementary elliptic integral is

$$B(w) = 1/\text{agm} \left(\sqrt{F(w)}, \sqrt{F(-w)} \right).$$

1.2 The equal-mass case

With $a = b = c = 1$, we have $F(w) = (w - 1)^3(w + 3)$ and hence

$$w^2 B(w) = G_3(1/w), \quad w^2 A(w) = \tilde{G}_3(1/w)$$

are given by complementary pair of **honeycomb** solutions

$$\begin{aligned} G_3(y) &= 1/\operatorname{agm}\left(\sqrt{(1-y)^3(1+3y)}, \sqrt{(1+y)^3(1-3y)}\right), \\ \tilde{G}_3(y) &= 1/\operatorname{agm}\left(\sqrt{(1-y)^3(1+3y)}, 4y\sqrt{y}\right). \end{aligned}$$

Defining the elliptic **nome** $q = \exp(-\pi B(w)/A(w))$, we have

$$-\left(q \frac{d}{dq}\right)^2 \left(\frac{I(w, 1, 1, 1)}{24\sqrt{3}A(w)}\right) = \frac{w^2(w^2 - 1)(w^2 - 9)A(w)^3}{9\sqrt{3}}$$

as the differential equation, found by Broadhurst, Fleischer and Tarasov.

Regarding w and $A(w)$ as functions of q , we obtain the **modular** functions

$$\begin{aligned} \frac{w}{3} &= \left(\frac{\eta_3}{\eta_1}\right)^4 \left(\frac{\eta_2}{\eta_6}\right)^2, \quad 4\sqrt{3}A = \frac{\eta_1^6 \eta_6}{\eta_2^3 \eta_3^2}, \\ \eta_n &= q^{n/24} \prod_{k>0} (1 - q^{nk}) = \sum_{k \in \mathbf{Z}} (-1)^k q^{n(6k+1)^2/24}. \end{aligned}$$

The two algebraic relations between $\{\eta_1, \eta_2, \eta_3, \eta_6\}$ give

$$\frac{w^2 - 1}{8} = \left(\frac{\eta_2}{\eta_1}\right)^9 \left(\frac{\eta_3}{\eta_6}\right)^3, \quad \frac{w^2 - 9}{72} = \left(\frac{\eta_6}{\eta_1}\right)^5 \frac{\eta_2}{\eta_3}.$$

Hence the BFT equation reduces to

$$-\left(q \frac{d}{dq}\right)^2 \left(\frac{I}{24\sqrt{3}A}\right) = \frac{w}{3} f_{3,12} = \left(\frac{\eta_3^3}{\eta_1}\right)^3 + \left(\frac{\eta_6^3}{\eta_2}\right)^3$$

where $f_{3,12} \equiv (\eta_2\eta_6)^3$ is a weight-3 level-12 modular form. Let $\chi(n) = \pm 1$ for $n = \pm 1 \pmod{6}$ and $\chi(n) = 0$ otherwise. Then

$$-\left(q \frac{d}{dq}\right)^2 \left(\frac{I}{24\sqrt{3}A}\right) = \sum_{n>0} \frac{n^2(q^n - q^{5n})}{1 - q^{6n}} = \sum_{n>0} \sum_{k>0} n^2 \chi(k) q^{nk}.$$

Integrating twice and using the known imaginary part on the cut, we get

$$\frac{I(w^2, 1, 1, 1)}{4A(w)} = E_2(q) = -\pi \log(-q) - 3\sqrt{3} \sum_{k>0} \frac{\chi(k)}{k^2} \frac{1 + q^k}{1 - q^k} = -E_2(1/q).$$

This **elliptic dilogarithm** was obtained by Bloch and Vanhove.

2 Diamond walks at 3 loops with Bessel

boob redder testes teammate reappear intestines interessierten

are **abelian squares**: words whose second halves are permutations of their first halves. There is a bijection between abelian squares of length $2k$ in an n -letter alphabet and **returning walks** of length $2k$ on the regular lattice in $n - 1$ dimensions with n -valent vertices. Thus the number $W_n(2k)$ of walks of length $2k$ in $n - 1$ dimensions is generated by

$$(I_0(2y))^n = \sum_{k=0}^{\infty} W_n(2k) \left(\frac{y^k}{k!}\right)^2$$

i.e. by the n -th power of the **Bessel** function $I_0(2y) = \sum_{k \geq 0} (y^k/k!)^2$.

Identity for diamond: It was shown by **Geoffrey Joyce** in 1973 that

$$G_4(z) = \sum_{k=0}^{\infty} W_4(2k)z^{2k} = (1 - y^2)(1 - 9y^2)G_3^2(y), \quad z^2 = \frac{-y^2}{(1 - y^2)(1 - 9y^2)}.$$

2.1 Three-loop sunrise with Bessel and Gauss

Joyce's transformation of $G_4(z)$ to the **square** of $G_3(y)$ enables progress with the equal-mass three-loop sunrise diagram

$$J(w) = 8 \int_0^\infty I_0(wt) K_0^4(t) t dt$$

using the transformation

$$\frac{w^2}{64} = \frac{-y^2}{(1-y^2)(1-9y^2)}$$

which is solved by

$$y = \frac{2}{\sqrt{4-w^2} + \sqrt{16-w^2}}$$

with singularities at the pseudo-threshold $w = 2$ and the physical threshold $w = 4$. Then the solutions to the third-order homogeneous equation for $J(w)$ are $(yG_3(y))^2$, $(y\tilde{G}_3(y))^2$ and $y^2G_3(y)\tilde{G}_3(y)$. Thus a strategy for best presenting the inhomogeneous equation is to divide J by one of these 3 and to operate with $(qd/dq)^3$, where $\log(q)$ is proportional to the ratio of a pair of solutions. If the result is expressible as a simple q series, then the problem may be solved in the same manner as at two loops.

Let $y = 2/(\sqrt{4 - w^2} + \sqrt{16 - w^2})$ and $q = \exp(-\frac{2}{3}\pi\tilde{G}_3(y)/G_3(y))$ with

$$G_3(y) = \frac{1}{\operatorname{agm}(\sqrt{(1-y)^3(1+3y)}, \sqrt{(1+y)^3(1-3y)})},$$

$$\tilde{G}_3(y) = \frac{1}{\operatorname{agm}(\sqrt{(1-y)^3(1+3y)}, 4y\sqrt{y})}.$$

Then the differential equation is

$$\left(q \frac{d}{dq}\right)^3 \left(\frac{2J(w)}{y^2 G_3^2(y)}\right) = -48 + 2 \sum_{n>0} \sum_{k>0} n^3 \psi(k) q^{nk}$$

with $\psi(k) = \psi(k+6) = \psi(6-k)$, and integers $\psi(1) = -48$, $\psi(2) = 720$, $\psi(3) = 384$, $\psi(6) = -5760$, that were found by Bloch, Kerr and Vanhove. We now integrate integrate 3 times. The constants of integration are determined by $J(0) = 7\zeta(3)$. The result is an **elliptic trilogarithm**:

$$\frac{2J(w)}{y^2 G_3^2(y)} = E_3(q) = (-2 \log(q))^3 + \sum_{k>0} \frac{\psi(k)}{k^3} \frac{1+q^k}{1-q^k} = -E_3(1/q).$$

3 Modular forms to 6 loops and a challenge for RISC

With $N = a + b$ **Bessel** functions and $c \geq 0$, I define **moments**

$$M(a, b, c) \equiv \int_0^\infty I_0^a(t) K_0^b(t) t^c dt$$

that converge for $b > a \geq 0$. For $b = a = N/2$, we have convergence for $b > c + 1$. The L -loop on-shell **sunrise** diagram in $D = 2$ dimensions gives

$$2^L M(1, L + 1, 1) = \int_0^\infty \cdots \int_0^\infty \frac{\prod_{k=1}^L dx_k/x_k}{(1 + \sum_{i=1}^L x_i)(1 + \sum_{j=1}^L 1/x_j) - 1}$$

as an integral over Schwinger parameters. $M(2, L, 1)$ is obtained by cutting an internal line. To obtain $M(1, L + 1, 3)$ and $M(2, L, 3)$, we differentiate w.r.t. an external momentum, before taking the **on-shell** limit.

Recently, in [arXiv:1706.08308], **Yajun Zhou** gave a complete **proof** of my 10-year-old conjecture on the 5-Bessel matrix:

$$\mathcal{M}_5 \equiv \begin{bmatrix} M(1, 4, 1) & M(1, 4, 3) \\ M(2, 3, 1) & M(2, 3, 3) \end{bmatrix} = \begin{bmatrix} \pi^2 C & \pi^2 \left(\frac{2}{15}\right)^2 \left(13C - \frac{1}{10C}\right) \\ \frac{\sqrt{15}\pi}{2} C & \frac{\sqrt{15}\pi}{2} \left(\frac{2}{15}\right)^2 \left(13C + \frac{1}{10C}\right) \end{bmatrix}.$$

The **determinant** $\det \mathcal{M}_5 = 2\pi^3/\sqrt{3^3 5^5}$ is **free** of the 3-loop constant

$$C \equiv \frac{\pi}{16} \left(1 - \frac{1}{\sqrt{5}}\right) \left(\sum_{n=-\infty}^{\infty} e^{-n^2 \pi \sqrt{15}}\right)^4 = \frac{1}{240\sqrt{5}\pi^2} \prod_{k=0}^3 \Gamma\left(\frac{2^k}{15}\right)$$

with Γ values from the **square** of an elliptic integral [arXiv:0801.0891] at the 15th singular value. The **L-series** for $N = 5$ Bessel functions comes from a **modular form** of weight 3 and level 15 [arXiv:1604.03057]:

$$\begin{aligned} \eta_n &\equiv q^{n/24} \prod_{k>0} (1 - q^{nk}) \\ f_{3,15} &\equiv (\eta_3 \eta_5)^3 + (\eta_1 \eta_{15})^3 = \sum_{n>0} A_5(n) q^n \\ L_5(s) &\equiv \sum_{n>0} \frac{A_5(n)}{n^s} \quad \text{for } s > 2 \\ \Lambda_5(s) &\equiv \left(\frac{15}{\pi^2}\right)^{s/2} \Gamma\left(\frac{s}{2}\right) \Gamma\left(\frac{s+1}{2}\right) L_5(s) = \Lambda_5(3-s) \\ L_5(1) &= \sum_{n>0} \frac{A_5(n)}{n} \left(2 + \frac{\sqrt{15}}{2\pi n}\right) \exp\left(-\frac{2\pi n}{\sqrt{15}}\right) \\ &= 5C = \frac{5}{\pi^2} \int_0^\infty I_0(t) K_0^4(t) t dt. \end{aligned}$$

3.1 Magnetic moment of the electron at $N = 6$

Here the **modular form**, found with **Francis Brown** in 2010, is

$$f_{4,6} \equiv (\eta_1 \eta_2 \eta_3 \eta_6)^2$$

with weight 4 and level 6. I discovered and **Zhou** proved that

$$2M(3, 3, 1) = 3L_6(2), \quad 2M(2, 4, 1) = 3L_6(3), \quad 2M(1, 5, 1) = \pi^2 L_6(2).$$

Stefano Laporta has evaluated 4-loop contributions to the **magnetic moment** of the **electron**. These engage the **first** row of the **determinant** [arXiv:1604.03057]

$$\det \begin{bmatrix} M(1, 5, 1) & M(1, 5, 3) \\ M(2, 4, 1) & M(2, 4, 3) \end{bmatrix} = \frac{5\zeta(4)}{32}.$$

It is notable that the hypergeometric series in

$$L_6(3) = \frac{\pi^2}{15} {}_4F_3 \left(\begin{matrix} \frac{1}{3}, & \frac{1}{2}, & \frac{1}{2}, & \frac{2}{3} \\ \frac{5}{6}, & 1, & \frac{7}{6} \end{matrix} \middle| 1 \right)$$

does not appear in Laporta's final result, though it was present at intermediate stages.

3.2 Kloosterman sums over finite fields

For $a \in \mathbf{F}_q$, with $q = p^k$, we define Kloosterman sums

$$K(a) \equiv \sum_{x \in \mathbf{F}_q^*} \exp\left(\frac{2\pi i}{p} \text{Trace}\left(x + \frac{a}{x}\right)\right)$$

with a trace of Frobenius in \mathbf{F}_q over \mathbf{F}_p . Then we obtain integers

$$c_N(q) \equiv -\frac{1 + S_N(q)}{q^2}, \quad S_N(q) \equiv \sum_{a \in \mathbf{F}_q^*} \sum_{k=0}^N [g(a)]^k [h(a)]^{N-k}$$

with $K(a) = -g(a) - h(a)$ and $g(a)h(a) = q$. Then

$$Z_N(p, T) = \exp\left(-\sum_{k>0} \frac{c_N(p^k)}{k} T^k\right)$$

is a polynomial in T . For $N < 8$ and $s > (N - 1)/2$, the L-series is

$$L_N(s) = \prod_p \frac{1}{Z_N(p, p^{-s})}.$$

With $N = 7$ Bessel functions, the **local** factors at the **primes** in

$$L_7(s) = \prod_p \frac{1}{Z_7(p, p^{-s})} \quad \text{for } s > 3$$

are given, for the **good** primes p coprime to 105, by the **cubic**

$$Z_7(p, T) = \left(1 - \left(\frac{p}{105}\right) p^2 T\right) \left(1 + \left(\frac{p}{105}\right) (2p^2 - |\lambda_p|^2) T + p^4 T^2\right)$$

where $\left(\frac{p}{105}\right) = \pm 1$ is a **Kronecker** symbol and λ_p is a Hecke eigenvalue of a weight-3 newform with level 525. For the primes of **bad** reduction, I obtained **quadratics** from **Kloosterman** moments in **finite fields**:

$$Z_7(3, T) = 1 - 10T + 3^4 T^2, \quad Z_7(5, T) = 1 - 5^4 T^2, \quad Z_7(7, T) = 1 + 70T + 7^4 T^2.$$

Then **Anton Mellit** suggested a **functional equation**

$$\Lambda_7(s) \equiv \left(\frac{105}{\pi^3}\right)^{s/2} \Gamma\left(\frac{s-1}{2}\right) \Gamma\left(\frac{s}{2}\right) \Gamma\left(\frac{s+1}{2}\right) L_7(s) = \Lambda_7(5-s)$$

that was validated at high precision and gave us the empirical result

$$24M(2, 5, 1) = 5\pi^2 L_7(2).$$

3.3 Subtleties at $N = 8$

With $N = 8$ Bessel functions, the L-series comes from the **modular form**

$$f_{6,6} \equiv \left(\frac{\eta_2^3 \eta_3^3}{\eta_1 \eta_6} \right)^3 + \left(\frac{\eta_1^3 \eta_6^3}{\eta_2 \eta_3} \right)^3$$

with weight 6 and level 6. I discovered and **Zhou** proved that

$$M(4, 4, 1) = L_8(3), \quad 4M(3, 5, 1) = 9L_8(4), \quad 4M(2, 6, 1) = 27L_8(5),$$

and $4M(1, 7, 1) = 9\pi^2 L_8(4)$ for the **6-loop sunrise** integral.

There are **two subtleties**. Kloosterman moments at $N = 8$ do **not** deliver the local factors directly: in $L_8(s) = \prod_p Z_4(p, p^{2-s})/Z_8(p, p^{-s})$ we remove factors from $N = 4$. Secondly, there is an infinite family of **sum rules**:

$$a(n) \equiv \left(\frac{2}{\pi} \right)^4 \int_0^\infty (\pi^2 I_0^2(t) - K_0^2(t)) I_0(t) K_0^5(t) (2t)^{2n-1} dt$$

delivers the **integers** of <http://oeis.org/A262961> as was recently proved by **Zhou** in [arXiv:1706.01068].

3.4 Eta-quotient challenge at weight 6 and level 24

Please verify and **simplify** these 3 newforms with weight 6 and level 24:

$$\begin{aligned}
E_{\pm} &:= \eta_3^4 \left(\left(\frac{\eta_4 \eta_6^3 \eta_8}{\eta_{24}} \right)^2 + 16 \left(\frac{\eta_{12} \eta_8^3 \eta_{24}}{\eta_6} \right)^2 - 48 \left(\frac{\eta_4 \eta_{24}^3 \eta_8}{\eta_6} \right)^2 \right) \\
&+ \eta_{24}^4 \left(\left(\frac{\eta_3 \eta_1^3 \eta_6}{\eta_{12}} \right)^2 + 3 \left(\frac{\eta_1 \eta_3^3 \eta_2}{\eta_{12}} \right)^2 - 16 \left(\frac{\eta_1 \eta_{12}^3 \eta_2}{\eta_3} \right)^2 \right) \\
&\pm \eta_8^4 \left(\left(\frac{\eta_3 \eta_1^3 \eta_6}{\eta_4} \right)^2 - \left(\frac{\eta_1 \eta_3^3 \eta_2}{\eta_4} \right)^2 - \frac{16}{3} \left(\frac{\eta_3 \eta_4^3 \eta_6}{\eta_1} \right)^2 \right) \\
&\pm \eta_1^4 \left(\frac{1}{3} \left(\frac{\eta_{12} \eta_2^3 \eta_{24}}{\eta_8} \right)^2 - 16 \left(\frac{\eta_{12} \eta_8^3 \eta_{24}}{\eta_2} \right)^2 - 16 \left(\frac{\eta_4 \eta_{24}^3 \eta_8}{\eta_2} \right)^2 \right) \\
F &:= \eta_3^4 \left(\frac{\eta_4 \eta_6^3 \eta_8}{\eta_{24}} \right)^2 + \frac{1}{3} \eta_1^4 \left(\frac{\eta_{12} \eta_2^3 \eta_{24}}{\eta_8} \right)^2 - 8 \eta_4^{12} - 216 \eta_{12}^{12} \\
&- \eta_{24}^4 \left(7 \left(\frac{\eta_3 \eta_1^3 \eta_6}{\eta_{12}} \right)^2 - 3 \left(\frac{\eta_1 \eta_3^3 \eta_2}{\eta_{12}} \right)^2 - 32 \left(\frac{\eta_1 \eta_{12}^3 \eta_2}{\eta_3} \right)^2 \right) \\
&+ \eta_8^4 \left(\left(\frac{\eta_3 \eta_1^3 \eta_6}{\eta_4} \right)^2 + 7 \left(\frac{\eta_1 \eta_3^3 \eta_2}{\eta_4} \right)^2 + \frac{32}{3} \left(\frac{\eta_3 \eta_4^3 \eta_6}{\eta_1} \right)^2 \right).
\end{aligned}$$

3.5 Vacuum integrals and non-critical modular L-series

In the **modular** cases $N = 5, 6, 8$, L-series **outside** the critical strip are empirically related to **determinants** that contain **vacuum** integrals:

$$\begin{aligned} \det \int_0^\infty K_0^3(t) \begin{bmatrix} K_0^2(t) & t^2 K_0^2(t) \\ I_0^2(t) & t^2 I_0^2(t) \end{bmatrix} t dt &= \frac{45}{8\pi^2} L_5(4) \\ \det \int_0^\infty K_0^4(t) \begin{bmatrix} K_0^2(t) & t^2 K_0^2(t) \\ I_0^2(t) & t^2 I_0^2(t) \end{bmatrix} t dt &= \frac{27}{4\pi^2} L_6(5) \\ \det \int_0^\infty K_0^6(t) \begin{bmatrix} K_0^2(t) & t^2(1-2t^2)K_0^2(t) \\ I_0^2(t) & t^2(1-2t^2)I_0^2(t) \end{bmatrix} t dt &= \frac{6075}{128\pi^2} L_8(7). \end{aligned}$$

3.6 Signpost

In work at $N > 8$ with **David Roberts** these features are notable:
local factors from **Kloosterman** moments, sometimes with adjustment;
guesses of Γ factors, signs and conductors in **functional equations**;
empirical fits of L-series to **determinants** of Feynman integrals;
quadratic relations between Bessel moments; **sum rules** when $4|N$.

4 L-series up to 22 loops

Let $\Omega_{a,b}$ be the **determinant** of the $r \times r$ matrix with $M(a, b, 1)$ at top left, size $r = \lceil (a + b)/4 - 1 \rceil$, powers of t^2 increasing to the right and powers of $I_0^2(t)$ increasing downwards. Thus $\Omega_{1,23}$ is a 5×5 determinant with the **22-loop sunrise** integral $M(1, 23, 1)$ at **top left** and $M(9, 15, 9)$ at bottom right. With $N = 4r + 4$ Bessel functions, we discovered that

$$\begin{aligned}
 L_8(4) &= \frac{2^2 \Omega_{1,7}}{3^2 \pi^2} \equiv \frac{4}{9\pi^2} \int_0^\infty I_0(t) K_0^7(t) t dt \\
 L_{12}(6) &= \frac{2^6 \Omega_{1,11}}{3^4 \times 5\pi^6} \\
 L_{16}(8) &= \frac{2^{14} \Omega_{1,15}}{3^7 \times 5^2 \times 7\pi^{12}} \\
 L_{20}(10) &= \frac{2^{22} \times 11 \times \mathbf{131} \Omega_{1,19}}{3^{11} \times 5^6 \times 7^3 \pi^{20}} \quad \text{to 44 digits} \\
 L_{24}(12) &= \frac{2^{29} \times \mathbf{12558877} \Omega_{1,23}}{3^{19} \times 5^9 \times 7^3 \times 11\pi^{30}} \quad \text{to 19 digits,}
 \end{aligned}$$

where boldface highlights **primes** greater than N . **30 GHz-years** of work gave 44-digit **precision** for $L_{20}(10)$. $L_{24}(12)$ agrees up to 19 digits.

With a **cut** of a line in the diagram at top left of the matrix, we found

$$\begin{aligned}
L_8(5) &= \frac{2^2 \Omega_{2,6}}{3^3} \equiv \frac{4}{27} \int_0^\infty I_0^2(t) K_0^6(t) t dt \\
L_{12}(7) &= \frac{2^5 \times 11 \Omega_{2,10}}{3^6 \times 5^2 \pi^2} \\
L_{16}(9) &= \frac{2^{14} \times 13 \Omega_{2,14}}{3^9 \times 5^3 \times 7^2 \pi^6} \\
L_{20}(11) &= \frac{2^{19} \times 17 \times 19 \times \mathbf{23} \Omega_{2,18}}{3^{13} \times 5^7 \times 7^3 \pi^{12}} \\
L_{24}(13) &= \frac{2^{27} \times 17 \times 19^2 \times 23^2 \times \mathbf{46681} \Omega_{2,22}}{3^{23} \times 5^{12} \times 7^4 \times 11^2 \pi^{20}}.
\end{aligned}$$

At $N = 12, 16, 20$, with an **odd** sign in the functional equation, we found

$$\begin{aligned}
-L'_{12}(5) &= \frac{2^4 (2^6 \times \mathbf{29} \widehat{\Omega}_{2,10} + 3 \Omega_{2,10} \log 2)}{3^2 \times 7 \pi^6} \\
-L'_{16}(7) &= \frac{2^9 (2^7 \times \mathbf{83} \widehat{\Omega}_{2,14} + 3 \times 11 \Omega_{2,14} \log 2)}{3^5 \times 5 \pi^{12}} \\
-L'_{20}(9) &= \frac{2^{17} \times 17 \times 19 (2^9 \times 7 \times \mathbf{101} \widehat{\Omega}_{2,18} + 5 \times 13 \Omega_{2,18} \log 2)}{3^8 \times 5^4 \times 7^2 \times 11 \pi^{20}}
\end{aligned}$$

for **central derivatives**, using **enlarged** determinants $\widehat{\Omega}_{2,4r+2}$ of size $r + 1$ with **regularization** of $M(2r + 2, 2r + 2, 2r + 1)$ at bottom right.

In the cases with $N = 4r + 2$, we obtained

$$\begin{aligned}
L_6(2) &= \frac{2\Omega_{1,5}}{\pi^2} \equiv \frac{2}{\pi^2} \int_0^\infty I_0(t)K_0^5(t)tdt \\
L_6(3) &= \frac{2\Omega_{2,4}}{3} \equiv \frac{2}{3} \int_0^\infty I_0^2(t)K_0^4(t)tdt \\
L_{10}(4) &= \frac{2^7\Omega_{1,9}}{3^2\pi^6} \\
L_{10}(5) &= \frac{2^4\Omega_{2,8}}{3 \times 5\pi^2} \\
L_{14}(6) &= 0 \\
L_{14}(7) &= \frac{2^{10} \times 11 \times 13 \Omega_{2,12}}{3^6 \times 5^2 \times 7\pi^6} \\
L_{18}(8) &= \frac{2^{21} \times 17 \times \mathbf{19} \Omega_{1,17}}{3^5 \times 5^4 \times 7\pi^{20}} \\
L_{18}(9) &= \frac{2^{12} \times 13 \times 17 \times \mathbf{41} \Omega_{2,16}}{3^8 \times 5^3 \times 7^2\pi^{12}} \\
L_{22}(10) &= 0 \\
L_{22}(11) &= \frac{2^{23} \times 17 \times 19 \times \mathbf{11621} \Omega_{2,20}}{3^{14} \times 5^7 \times 7^3\pi^{20}}
\end{aligned}$$

with central vanishing from an odd sign at $N = 14$ and $N = 22$.

For cases with odd N , we obtained

$$\begin{aligned}
L_5(2) &= \frac{2^2 \Omega_{2,3}}{3} \equiv \frac{4}{3} \int_0^\infty I_0^2(t) K_0^3(t) t dt \\
L_7(2) &= \frac{2^3 \times 3 \Omega_{2,5}}{5\pi^2} \equiv \frac{24}{5\pi^2} \int_0^\infty I_0^2(t) K_0^5(t) t dt \\
L_9(4) &= \frac{2^6 \Omega_{2,7}}{3 \times 5\pi^2} \\
L_{11}(4) &= \frac{2^8 \times 5 \Omega_{2,9}}{3 \times 7\pi^6} \\
L_{13}(6) &= \frac{2^7 \times \mathbf{149} \Omega_{2,11}}{3^3 \times 5 \times 7\pi^6} \\
L_{15}(6) &= \frac{2^8 \times 7 \times \mathbf{53} \Omega_{2,13}}{3^2 \times 5\pi^{12}} && \text{to 43 digits} \\
L_{17}(8) &= \frac{2^{15} \times \mathbf{29} \Omega_{2,15}}{3^5 \times 5^2 \times 7\pi^{12}} && \text{to 23 digits} \\
L_{19}(8) &= \frac{2^{14} \times \mathbf{1093} \times \mathbf{13171} \Omega_{2,17}}{3^4 \times 5^4 \times 7 \times 11\pi^{20}} && \text{to 14 digits.}
\end{aligned}$$

Comment: We also have results relating Bessel moments $M(a, b, c)$ with **even** c to L-series from Kloosterman moments with a quadratic **twist**.

5 Betti and de Rham matrices for all loops

Construction: Let v_k and w_k be the rational numbers **generated** by

$$\begin{aligned}\frac{J_0^2(t)}{C(t)} &= \sum_{k \geq 0} \frac{v_k}{k!} \left(\frac{t}{2}\right)^{2k} = 1 - \frac{17t^2}{54} + \frac{3781t^4}{186624} + \dots \\ \frac{2J_0(t)J_1(t)}{tC(t)} &= \sum_{k \geq 0} \frac{w_k}{k!} \left(\frac{t}{2}\right)^{2k} = 1 - \frac{41t^2}{216} + \frac{325t^4}{186624} + \dots\end{aligned}$$

where $J_0(t) = I_0(it)$, $J_1(t) = -J_0'(t)$ and

$$C(t) \equiv \frac{32(1 - J_0^2(t) - tJ_0(t)J_1(t))}{3t^4} = 1 - \frac{5t^2}{27} + \frac{35t^4}{2304} - \frac{7t^6}{9600} + \dots$$

We construct rational bivariate polynomials by the **recursion**

$$\begin{aligned}H_s(y, z) &= zH_{s-1}(y, z) - (s-1)yH_{s-2}(y, z) \\ &\quad - \sum_{k=1}^{s-1} \binom{s-1}{k} (v_k H_{s-k}(y, z) - w_k z H_{s-k-1}(y, z))\end{aligned}$$

for $s > 0$, with $H_0(y, z) = 1$. We use these to define

$$d_s(N, c) \equiv \frac{H_s(3c/2, N+2-2c)}{4^s s!}.$$

Matrices: We construct rational **deRham** matrices, with elements

$$D_N(a, b) \equiv \sum_{c=-b}^a d_{a-c}(N, -c)d_{b+c}(N, c)c^{N+1}$$

and a and b running from 1 to $k = \lceil N/2 - 1 \rceil$.

We act on those, on the left, with **period** matrices whose elements are

$$P_{2k+1}(u, a) \equiv \frac{(-1)^{a-1}}{\pi^u} M(k+1-u, k+u, 2a-1)$$

$$P_{2k+2}(u, a) \equiv \frac{(-1)^{a-1}}{\pi^{u+1/2}} M(k+1-u, k+1+u, 2a-1)$$

and on the right with their **transposes**, to define **Betti** matrices

$$B_N \equiv P_N D_N P_N^{\text{tr}}.$$

Conjecture: The Betti matrices have **rational** elements given by

$$B_{2k+1}(u, v) = (-1)^{u+k} 2^{-2k-2} (k+u)! (k+v)! Z(u+v)$$

$$B_{2k+2}(u, v) = (-1)^{u+k} 2^{-2k-3} (k+u+1)! (k+v+1)! Z(u+v+1)$$

$$Z(m) = \frac{1 + (-1)^m}{(2\pi)^m} \zeta(m).$$

Summary

1. Moments of 4 Bessel functions relate to walks on a **honeycomb**.
2. Moments of 5 Bessel functions relate to walks in a **diamond** crystal.
3. For $N = 5, 6$ and 8 Bessel functions, the **L-series** are **modular**.
4. For $N = 7$ and $N > 8$, **Kloosterman** moments yield local factors.
5. Relations between **determinants** of Feynman integrals and **L-series** have been discovered up to 22 loops and presumably go on for **ever**.
6. There are **quadratic** relations of the form $P_N D_N P_N^{\text{tr}} = B_N$ with **period, de Rham** and **Betti** matrices that we have specified.
7. Parallel results for **even** moments lead to an **eta-quotient challenge** for Günter Köhler, Peter Paule, Carsten Schneider, et alia.

I thank kind hosts in Creswick (Victoria), Newcastle (NSW), Mainz, Oxford, Paris, Marseille, Edinburgh, Copenhagen, Vienna and Zeuthen.