

Two-loop conformal anomaly in QCD

A. N. MANASHOV

Hamburg University

based on

V. Braun, A. Manashov, Eur. Phys. J. C **73** (2013) 2544

V. Braun, A. Manashov, Phys. Lett. B **734** (2014) 137

V. Braun, A. Manashov, S. Moch, M. Strohmaier, in progress

DESY, 1.10.2015

- **DIS: Structure functions** : scale dependence is governed by anomalous dimensions of twist-2 operators (known at three loops), **Larin, Vermaseren, Moch, Vogt, et al**)
- **Deeply Virtual Compton Scattering:** (**Müller, Ji, Radyushkin**) Generalized Parton Distributions: scale dependence \longleftrightarrow anomalous dimension matrix (nonforward kernel) (two loops) **Belitsky, Müller, (2000)**

Explore the road to NNLO (three-loop) evolution equations for GPDs

One loop: anomalous dimensions+conformal symmetry → full anomalous dimension matrix.
(Makeenko, 1980)

$$O_N \sim (\partial_{z_1} + \partial_{z_2})^N C_N^{3/2} \left(\frac{\partial_{z_1} - \partial_{z_2}}{\partial_{z_1} + \partial_{z_2}} \right) \bar{q}(z_1 n) \gamma_+ q(z_2 n)$$

In any realistic $d = 4$ QFT the conformal symmetry is broken, $\beta(g) \neq 0$.

D. Müller, Constraints for anomalous dimensions of local light cone operators in ϕ^3 in six-dimensions theory, Z. Phys. C 49 (1991) 293.

(Conformal Ward Identities, Conformal anomaly, Conformal scheme, etc)

Belitsky, Müller, (2000) two loop kernels in QCD.

Difference to D.Müller:

Instead of considering consequences of broken conformal symmetry in QCD we make use of exact conformal symmetry of a modified theory: Large N_f QCD in $4 - 2\epsilon$ dimensions at critical coupling: Banks, Zaks, 82

$$\beta^{QCD}(a) = 2a[-\epsilon - \beta_0 a + \dots] \quad a_* = -4\pi\epsilon/\beta_0 + \dots \quad \beta^{QCD}(a_*) = 0$$


$$a_* = a_*(\epsilon) \quad \longleftrightarrow \quad \epsilon = \epsilon(a_*)$$

$1/N_f$ expansion, Gracey, Cuccini, Derkachov, A.M.

Premium:

- Exact symmetry \Rightarrow algebraic group-theory methods
- Answer obtained directly in $\overline{\text{MS}}$ scheme
- Light-ray operator basis, do not need to restore evolution eqs. from local operators.

$$\begin{aligned} \mathcal{O}(x; z_1, z_2) &\equiv [\bar{q}(x + z_1 n) \not{n} q(x + z_2 n)] \equiv \sum_{m,k} \frac{z_1^m z_2^k}{m!k!} [(D_+^m \bar{q})(x) \not{n} (D_+^k q)(x)] \\ &= \sum_{N,k} \Psi_{Nk}(z_1, z_2) \partial_+^k \mathcal{O}_N(x) \end{aligned}$$



$$\begin{aligned}
 p_+ &= \frac{1}{\sqrt{2}}(p_0 + p_z) \rightarrow \infty \\
 p_- &= \frac{1}{\sqrt{2}}(p_0 - p_z) \rightarrow 0 \\
 px &\rightarrow p_+ x_-
 \end{aligned}$$

- Special conformal transformation

$$x_- \rightarrow x'_- = \frac{x_-}{1 + 2ax_-}$$

- translations $x_- \rightarrow x'_- = x_- + c$
- dilatations $x_- \rightarrow x'_- = \lambda x_-$

form the so-called **collinear subgroup** $SL(2, R)$

$$\alpha \rightarrow \alpha' = \frac{a\alpha + b}{c\alpha + d}, \quad ad - bc = 1$$

$$\Phi(\alpha) \rightarrow \Phi'(\alpha) = (c\alpha + d)^{-2j} \Phi\left(\frac{a\alpha + b}{c\alpha + d}\right)$$

where $\Phi(x) \rightarrow \Phi(x_-) = \Phi(\alpha n_-)$ is the quantum field with scaling dimension ℓ and spin projection s “living” on the light-ray

Conformal spin:

$$j = (\ell + s)/2$$

Light-ray operators satisfy the RG equation

Balitsky, Braun '89

$$\left(M\partial_M + \beta(a)\partial_a + \mathbb{H} \right) \mathcal{O}(z_1, z_2) = 0$$

where \mathbb{H} is an integral operator acting on the light-cone coordinates of the fields:

$$\mathbb{H}\mathcal{O}(z_1, z_2) = \int_0^1 d\alpha \int_0^1 d\beta h(\alpha, \beta) \mathcal{O}(z_{12}^\alpha, z_{21}^\beta)$$

$$\begin{aligned} z_{12}^\alpha &\equiv z_1 \bar{\alpha} + z_2 \alpha \\ \bar{\alpha} &= 1 - \alpha \end{aligned}$$

$$h(\alpha, \beta) = a h^{(1)}(\alpha, \beta) + a^2 h^{(2)}(\alpha, \beta) + \dots \text{ [does not depend on } \epsilon \text{]}$$

One can show that the powers $\mathcal{O}(z_1, z_2) \mapsto (z_1 - z_2)^N$ are eigenfunctions of \mathbb{H} , and the corresponding eigenvalues are the anomalous dimensions of local operators of spin N (with $N - 1$ derivatives)

$$\gamma_N = \int d\alpha d\beta h(\alpha, \beta) (1 - \alpha - \beta)^{N-1} \quad \text{(NNLO)}$$

Is it possible to restore $h(\alpha, \beta)$ from γ_N ?

It is expected that the evolution kernel \mathbb{H} commutes with the generators of $SL(2, R)$ collinear subgroup, $[S_\alpha, \mathbb{H}] = 0$, $\alpha = \pm, 0$.

- Leading order generators

$$S_+^{(0)} = z_1^2 \partial_{z_1} + z_2^2 \partial_{z_2} + 2(z_1 + z_2), \quad S_0^{(0)} = z_1 \partial_{z_1} + z_2 \partial_{z_2} + 2, \quad S_-^{(0)} = -\partial_{z_1} - \partial_{z_2}$$

- Exact conformal symmetry, but the generators are modified by quantum corrections

$$S_- = S_-^{(0)},$$

$$S_0 = S_0^{(0)} - \epsilon(a_*) + \frac{1}{2} \mathbb{H}(a_*), \quad \mathbb{H}(a_*) = a_* \mathbb{H}^{(1)} + \dots$$

$$S_+ = S_+^{(0)} + (z_1 + z_2) \left(-\epsilon(a_*) + \frac{1}{2} \mathbb{H}(a_*) \right) + (z_1 - z_2) \Delta_+(a_*),$$

$$a = \frac{\alpha_s}{4\pi}$$

$$\mathcal{O}(x; z_1, z_2) = \sum_{N,k} \Psi_{Nk}(z_1, z_2) \partial_+^k \mathcal{O}_N(x), \quad \mathbb{H}(a_*) \Psi_{Nk} = \gamma_N \Psi_{Nk}$$

$\mathbb{H}(a_*)$ commutes with all generators $\implies \mathbb{H}(a_*) = f(J)$

$$J(J-1) = S_+ S_- + S_0(S_0-1)$$

The lowest eigenfunctions are known: $\Psi_{N0}(z_1, z_2) = z_{12}^N$ (for any coupling).

$$J(a_*) z_{12}^N = (N+2-\epsilon + \gamma_N/2) z_{12}^N, \quad \mathbb{H}(a_*) z_{12}^N = \gamma_N z_{12}^N$$

$$\gamma_N = f(N+2-\epsilon + \gamma_N/2).$$

One can restore the function f from anomalous dimensions: $f(x) = a f_1(x) + a^2 f_2(x) + \dots$

$$\mathbb{H}(a) = a f_1(J(a)) + a^2 f_2(J(a)) + a^3 f_3(J(a)) + \dots$$

- Expansion over conformal operators

$$\mathcal{O}^{(n)}(x; z_1, z_2) = \sum_{N,k} \Psi_{Nk}(z_1, z_2) \partial_+^k \mathcal{O}_N^{(n)}(x),$$

- Conformal symmetry at the critical coupling implies

$$\left(S_+^{(z)} - \frac{1}{2} x^2 (\bar{n} \partial_x) \right) \langle \mathcal{O}_n(0, z_1, z_2), \mathcal{O}_{\bar{n}}(x, w_1, w_2) \rangle = 0 \quad \boxed{(nx) = (\bar{n}x) = 0}$$

- To find explicit expression for S_+ , consider Ward identity ($\delta_+ = \mathbf{K} \bar{n}$)

$$\langle \delta_+ \mathcal{O}^{(n)}(z) \mathcal{O}^{(\bar{n})}(x, w) \rangle + \langle \mathcal{O}^{(n)}(z) \delta_+ \mathcal{O}^{(\bar{n})}(x, w) \rangle = \langle \delta_+ S_R \mathcal{O}^{(n)}(z) \mathcal{O}^{(\bar{n})}(x, w) \rangle$$

Then

$$\delta_+ \mathcal{O}^{(\bar{n})}(x; w_1, w_2) = -x^2 (\bar{n} \partial_x) \mathcal{O}^{(\bar{n})}(x; w_1, w_2)$$

$$\delta_+ \mathcal{O}^{(n)}(0; z_1, z_2) = 2(n\bar{n}) \left(S_+^{(0)} - \epsilon(z_1 + z_2) - \frac{a}{2} [\mathbb{H}^{(1)}, z_1 + z_2] + \dots \right) \mathcal{O}^{(n)}(0; z_1, z_2)$$

and for the last term

$$\delta_+ S^{QCD} = 4\epsilon \int d^d x (x \bar{n}) L^{QCD} + 2(d-2) \bar{n}^\mu \int d^d x \delta_{BRST} (\bar{c}^a A_\mu^a).$$

$$\langle \mathcal{O}^{(n)}(0; z_1, z_2) q(x) \bar{q}(y) \rangle - \text{bad object for an analysis}$$

- Reexpand ϵL in terms of renormalized operators

$$2\epsilon L = -\beta(a)/a [L^{YM+gf}] + EOM + BRST$$

$$[\beta(a_*) = 0]$$

$$\Delta S_+ \langle \mathcal{O}_n(0, z), \mathcal{O}_{\bar{n}}(x, w) \rangle = KR' \left(\left\langle \int d^D y (\bar{n}y) L^{YM+gf}(y) \mathcal{O}_n(0, z), \mathcal{O}_{\bar{n}}(x, w) \right\rangle \right)$$

SIMPLE RESIDUE IN ϵ

One loop: **A. Belitsky, D. Müller, 2001, (V. Braun, A.M., (2014))**

$$\Delta_+^{(1)} \mathcal{O}(z_1, z_2) = -2C_F \int_0^1 du \int_0^1 d\alpha \frac{\bar{\alpha}}{\alpha} \left[\mathcal{O}(z_{12}^{u\alpha}, z_2) - \mathcal{O}(z_1, z_{21}^{u\alpha}) \right]$$

two loops:

$$\begin{aligned}
 [\Delta_+^{(2)} \mathcal{O}](z_1, z_2) &= \int_0^1 d\alpha \int_0^{\bar{\alpha}} d\beta \left[\omega(\alpha, \beta) + \omega^{\mathbb{P}}(\alpha, \beta) \mathbb{P}_{12} \right] \left[\mathcal{O}(z_{12}^\alpha, z_{21}^\beta) - \mathcal{O}(z_{12}^\beta, z_{21}^\alpha) \right] \\
 &\quad + \int_0^1 du \int_0^1 d\alpha \varkappa(\alpha) \left[\mathcal{O}(z_{12}^{u\alpha}, z_2) - \mathcal{O}(z_1, z_{21}^{u\alpha}) \right].
 \end{aligned}$$

Color structures: $\beta_0 C_F$, C_F^2 and $C_F C_A$: ($\tau = \alpha\beta/\bar{\alpha}\bar{\beta}$)

$$\begin{aligned}
 \omega_{FF}(\alpha, \beta) &= 4 \left[\left(\alpha - \frac{1}{\alpha} \right) \left[\text{Li}_2 \left(\frac{\beta}{\bar{\alpha}} \right) - \text{Li}_2(\beta) - \text{Li}_2(\alpha) - \frac{1}{4} \ln^2 \bar{\alpha} \right] - \alpha \left[\text{Li}_2(\alpha) - \text{Li}_2(1) \right] \right. \\
 &\quad \left. - \frac{\alpha + \beta}{2} \ln \alpha \ln \bar{\alpha} + \frac{1}{4} \left[\beta \ln^2 \bar{\alpha} - \alpha \ln^2 \alpha \right] - \frac{\alpha}{\tau} \left[\tau \ln \tau + \bar{\tau} \ln \bar{\tau} \right] \right. \\
 &\quad \left. + \frac{1}{4} \left[\beta - 2\bar{\alpha} + \frac{2\beta}{\alpha} \right] \ln \bar{\alpha} + \frac{1}{2} \left[\bar{\alpha} - \frac{\alpha}{\bar{\alpha}} - 3\beta \right] \ln \alpha - \frac{15}{4} \alpha \right],
 \end{aligned}$$

$$\begin{aligned}
 \omega_{FA}(\alpha, \beta) &= 2 \left[\left(\frac{1}{\alpha} - \alpha \right) \left[\text{Li}_2 \left(\frac{\beta}{\bar{\alpha}} \right) - \text{Li}_2(\beta) - 2 \text{Li}_2(\alpha) - \ln \alpha \ln \bar{\alpha} \right] + \frac{\alpha}{\tau} \left[\tau \ln \tau + \bar{\tau} \ln \bar{\tau} \right] \right. \\
 &\quad \left. - \bar{\beta} \ln \alpha - \frac{\bar{\alpha}}{\alpha} \ln \bar{\alpha} \right],
 \end{aligned}$$

$$\omega_{FF}^{\mathbb{P}}(\alpha, \beta) = -4 \left[\left(\bar{\alpha} - \frac{1}{\bar{\alpha}} \right) \left[\text{Li}_2 \left(\frac{\alpha}{\bar{\beta}} \right) - \text{Li}_2(\alpha) - \ln \bar{\alpha} \ln \bar{\beta} \right] + \alpha \bar{\tau} \ln \bar{\tau} + \frac{\beta^2}{\bar{\beta}} \ln \bar{\alpha} \right]$$

- QCD evolution equations possess a "hidden" conformal symmetry
- n -loop evolution kernels for twist-two operators in $\overline{\text{MS}}$ scheme can be restored from the $(n - 1)$ -loop calculation of the special conformal anomaly and n -loop anomalous dimensions
- $SL(2)$ symmetry properties manifest in the light-ray operator representation