

## Detection of reactor neutrinos with a delayed signal of neutron capture on Hydrogen at RENO

### Authorship annotation

for the RENO collaboration

### Session and Location

Monday Session, Poster Wall #161 (Ballroom)

### Abstract content

The RENO experiment has been taking electron antineutrino data from the reactors at Hanbit nuclear power plant in Korea, using two identical detectors since 2011. It has measured the neutrino mixing angle  $\theta_{13}$  using inverse beta decay events with a neutron captured on Gadolinium(Gd). A neutron is also captured by Hydrogen. Reactor antineutrinos are detected with a delayed signal of neutron capture by Hydrogen (n-H). The n-H sample provides an independent result and a consistent check of the n-Gd result. The coincidence time between the prompt and delayed events is longer than that of n-Gd, and results in a larger accidental background. We measure background rates in the 2000 day n-H data sample. In this presentation, we report the n-H selection criteria for reactor antineutrino candidates and remained background rates.

### Poster included in proceedings:

yes

**Primary author(s) :** Dr. KWON, Eunhyang (Seoul National University)

**Presenter(s) :** Mr. SHIN, ChangDong (Chonnam National University)

**Session Classification :** Poster Session Monday

**Track Classification :** Poster (not participating in poster prize competition)