

## Response of the STEREO detector

### Authorship annotation

on behalf of STEREO collaboration

### Session and Location

Monday Session, Poster Wall #195 (Ballroom)

### Abstract content

The STEREO experiment searches for light sterile neutrinos in the eV range. The STEREO detector is divided in two subvolumes both filled with liquid scintillator: an inner neutrino target loaded with gadolinium, which is subsequently subdivided in 6 different cells, and an external unloaded gamma catcher subdivided in 4 cells.

The response of each of the cells, including charge to energy conversion and energy non linearities, have been characterized by means of a set of gamma and neutron sources covering a broad range of the energy spectrum of reactor antineutrinos. In addition, the particular response of single photomultiplier tubes and the linearity of electronics are controlled hourly by a system of LEDs performing single photoelectron calibration.

This poster will discuss this characterization as a means to obtain an accurate control energy scale and detection efficiency of STEREO, together with the monitoring of the detector's stability and light cross-talk between cells.

### Poster included in proceedings:

yes

**Primary author(s) :** Mr. ROCA, Christian (Max-Planck-Institut für Kernphysik); Ms. ALMAZÁN, Helena (Max-Planck-Institut für Kernphysik)

**Co-author(s) :** Dr. SERGEYEVA, Viktoriya (Laboratoire d'Annecy de Physique des Particules, CNRS)

**Presenter(s) :** Mr. ROCA, Christian (Max-Planck-Institut für Kernphysik); Ms. ALMAZÁN, Helena (Max-Planck-Institut für Kernphysik)

**Session Classification :** Poster Session Monday

**Track Classification :** Poster (participating in poster prize competition)